

PARTIAL DIFFERENTIAL EQUATIONS

XAVIER ROS OTON

1. OVERVIEW AND PRELIMINARIES

- (1) Let $f : \Omega \subset \mathbb{C} \rightarrow \mathbb{C}$ be any holomorphic function, and let $u := \operatorname{Re} f$ and $v = \operatorname{Im} f$ be the real and imaginary parts of f .

Prove that, if we identify $\mathbb{C} \simeq \mathbb{R}^2$, they satisfy $\Delta u = 0$ and $\Delta v = 0$ in $\Omega \subset \mathbb{R}^2$.

(2 points)

- (2) Assume $\vec{\mathbf{E}}$ and $\vec{\mathbf{B}}$ solve Maxwell's equations in \mathbb{R}^3

$$\partial_t \vec{\mathbf{E}} = \operatorname{curl} \vec{\mathbf{B}}$$

$$\partial_t \vec{\mathbf{B}} = -\operatorname{curl} \vec{\mathbf{E}}$$

$$\operatorname{div} \vec{\mathbf{B}} = \operatorname{div} \vec{\mathbf{E}} = 0.$$

Prove that

$$\partial_{tt} \vec{\mathbf{E}} - \Delta \vec{\mathbf{E}} = 0 \quad \text{and} \quad \partial_{tt} \vec{\mathbf{B}} - \Delta \vec{\mathbf{B}} = 0.$$

(2 points)

- (3) Prove that, for any radial function $u \in C^2(\mathbb{R}^n)$, we have

$$\Delta u = \partial_{rr} u + \frac{n-1}{r} \partial_r u = r^{1-n} \partial_r (r^{n-1} \partial_r u),$$

where $r = |x|$.

(3 points)

- (4) Let $f \in C^\infty(\mathbb{R}^n)$. Prove, for $k = 1, 2, \dots$, the following Taylor expansion

$$f(x) = \sum_{|\alpha| \leq k} \frac{1}{\alpha!} D^\alpha f(0) x^\alpha + O(|x|^{k+1}) \quad \text{as } x \rightarrow 0,$$

where the sum is taken over multiindices $\alpha \in \mathbb{N}^n$.

Hint: Fix $x \in \mathbb{R}^n$ and consider the function of one variable $g(t) := f(tx)$.

(2 points)

- (5) Show that for any C^2 function f we have

$$\int_{B_r(x_0)} f - f(x_0) = c_n r^2 \Delta f(x_0) + o(r^2),$$

where $f_E f = \frac{1}{|E|} \int_E f$ is the average of f over the set E , and $c_n > 0$ is a constant.

Hint: Use Taylor's expansion.

(3 points)

(6) Show that for any C^1 vector field $\vec{F} : \mathbb{R}^n \rightarrow \mathbb{R}^n$ we have

$$\operatorname{div} \vec{F}(x) = \frac{1}{|B_1|} \lim_{r \rightarrow 0} \int_{\partial B_1} \vec{F}(x + r\theta) \cdot \theta \, dS(\theta).$$

(2 points)

(7) Let $\Omega \subset \mathbb{R}^n$ be any bounded smooth domain. Deduce the following integration by parts formulas from the divergence theorem: For any $f, g \in C^1(\overline{\Omega})$ we have

$$\int_{\Omega} f \partial_{x_i} g = - \int_{\Omega} g \partial_{x_i} f + \int_{\partial\Omega} f g \nu_i,$$

where ν_i is the i -th component of the normal vector ν . In particular, for any $u, w \in C^2(\overline{\Omega})$,

$$\int_{\Omega} \nabla w \cdot \nabla u = - \int_{\Omega} w \Delta u + \int_{\partial\Omega} w \nabla u \cdot \nu.$$

(3 points)

(8) Let $\Omega \subset \mathbb{R}^n$ be a bounded smooth domain. Integrate by parts to prove the interpolation inequality

$$\|\nabla u\|_{L^2(\Omega)}^2 \leq C \|u\|_{L^2(\Omega)} \|D^2 u\|_{L^2(\Omega)}$$

for all $u \in C_c^\infty(\Omega)$.

(2 points)

(9) Let $\Omega \subset \mathbb{R}^n$ be a bounded smooth domain. Prove the *trace inequality*

$$\int_{\partial\Omega} |u|^2 \leq C \left(\int_{\Omega} |\nabla u|^2 + |u|^2 \right)$$

for all $u \in C^\infty(\overline{\Omega})$.

Hint: Use that there exists a smooth vector field \vec{X} such that $\vec{X} \cdot \nu \geq 1$ on $\partial\Omega$. Then, apply the divergence theorem to $\int_{\partial\Omega} u^2 \vec{X} \cdot \nu$ to prove the result.

(2 points)

(10) Let $\Omega \subset \mathbb{R}^n$ be a bounded smooth domain.

(i) Use the previous exercise to show that there is a bounded linear operator

$$\operatorname{Tr} : H^1(\Omega) \rightarrow L^2(\partial\Omega)$$

such that $\operatorname{Tr} u = u|_{\partial\Omega}$ for any $u \in C^\infty(\overline{\Omega})$. We call it the *trace operator*.

(ii) Prove that there does not exist a bounded linear operator

$$\operatorname{Tr} : L^2(\Omega) \rightarrow L^2(\partial\Omega)$$

such that $\operatorname{Tr} u = u|_{\partial\Omega}$ for any $u \in C^\infty(\overline{\Omega})$.

Note: This means that we cannot talk about the boundary values of a function in $L^2(\Omega)$, however all functions in $H^1(\Omega)$ do have boundary values in $L^2(\partial\Omega)$.

(4 points)

- (11) Prove the Poincaré inequality in dimension $n = 1$, that is:

$$\int_a^b u^2 \leq C_{a,b} \int_a^b |u'|^2 \quad \text{if} \quad u(a) = u(b) = 0$$

for some constant $C_{a,b}$ depending only on a and b .

(3 points)

- (12) Let $\Omega \subset \mathbb{R}^n$ be a bounded smooth domain. Use Rellich compactness theorem to prove the following Poincaré inequality

$$\|u - \bar{u}_\Omega\|_{L^2(\Omega)} \leq C \|\nabla u\|_{L^2(\Omega)} \quad \text{for all } u \in H^1(\Omega),$$

where $\bar{u}_\Omega = \frac{1}{|\Omega|} \int_\Omega u$ is the average value of u in Ω , and C depends only on Ω and n .

Note: We do not assume $u = 0$ on $\partial\Omega$ here.

(4 points)

- (13) Prove that for any $u \in C_c^\infty(\mathbb{R}^n)$, we have

$$\sup_{\mathbb{R}^n} |u| \leq \|D^n u\|_{L^1(\mathbb{R}^n)}$$

where $D^n u$ denotes the n -th derivatives of u .

Hint: Use the Fundamental Theorem of Calculus n times, with respect to the variables x_1, x_2, \dots, x_n .

(2 points)