


WIGGLYHEDRA

ASILATA BAPAT AND VINCENT PILAUD

ABSTRACT. Motivated by categorical representation theory, we define the wiggly complex, whose vertices are arcs wiggling around $n + 2$ points on a line, and whose faces are sets of wiggly arcs which are pairwise pointed and non-crossing. The wiggly complex is a $(2n - 1)$ -dimensional pseudomanifold, whose facets are wiggly pseudotriangulations. We show that wiggly pseudotriangulations are in bijection with wiggly permutations, which are permutations of $[2n]$ avoiding the patterns $(2j - 1) \cdots i \cdots (2j)$ for $i < 2j - 1$ and $(2j) \cdots k \cdots (2j - 1)$ for $k > 2j$. These permutations define the wiggly lattice, an induced sublattice of the weak order. We then prove that the wiggly complex is isomorphic to the boundary complex of the polar of the wigglyhedron, for which we give explicit and simple vertex and facet descriptions. Interestingly, we observe that any Cambrian associahedron is normally equivalent to a well-chosen face of the wigglyhedron. Finally, we recall the correspondence of wiggly arcs with objects in a category, and we develop categorical criteria for a subset of wiggly arcs to form a face of the wiggly complex.



AB was partially supported by the Australian Research Council DECRA grant DE240100447. VP was partially supported by the Spanish grant PID2022-137283NB-C21 of MCIN/AEI/10.13039/501100011033 / FEDER, UE, by Departament de Recerca i Universitats de la Generalitat de Catalunya (2021 SGR 00697), by the French grant CHARMS (ANR-19-CE40-0017), and by the French–Austrian projects PAGCAP (ANR-21-CE48-0020 & FWF I 5788).

CONTENTS

1. Introduction	2
2. Wiggly pseudotriangulations and wiggly permutations	4
2.1. Wiggly pseudotriangulations and the wiggly complex	4
2.2. Wiggly permutations and the wiggly lattice	11
2.3. Bijection	13
2.4. Graph properties	15
3. Wiggly fan and wigglyhedron	16
3.1. Polyhedral geometry	16
3.2. \mathbf{g} -vectors and \mathbf{c} -vectors	17
3.3. Wiggly fan	18
3.4. Wigglyhedron	20
4. Further topics	22
4.1. Cambrian considerations	22
4.2. Wiggly pseudotriangulations of planar point sets	25
5. Categorical interpretation of wiggly pseudotriangulations	26
5.1. A triangulated category associated to a quiver	26
5.2. The Khovanov–Seidel curve model	28
5.3. Cohomology filtrations of spherical objects	30
5.4. Categorical constraints on compatibility of curves	31
Acknowledgements	32
References	32

1. INTRODUCTION

In this paper we consider wiggly pseudotriangulations, which are built from arcs wiggling around points on a line. We study their combinatorics in detail, construct polyhedral realizations of their flip graph, and interpret them in terms of categorical representation theory.

Combinatorics. Fix $n \geq 1$ and consider the $n+2$ points on the horizontal axis labelled $0, \dots, n+1$. A *wiggly arc* is an abscissa monotone curve starting at some point i , ending at some point j (with $0 \leq i < j \leq n+1$), and passing above or below each point $i+1, \dots, j-1$. Two wiggly arcs are *compatible* if they are pointed (the left endpoint of one is not the right endpoint of the other) and non-crossing (the interiors of the curves are disjoint). A *wiggly dissection* is a collection of pairwise compatible wiggly arcs, and a *wiggly pseudotriangulation* is an inclusion maximal wiggly dissection. The *wiggly complex* WC_n is the simplicial complex of wiggly dissections (minus their external wiggly arcs). We first establish the following result.

Proposition. *The wiggly complex WC_n is a $(2n-1)$ -dimensional pseudomanifold without boundary.*

In particular, all wiggly pseudotriangulations have $2n-1$ internal wiggly arcs, each of which can be *flipped*. We actually describe the flip in terms of wiggly diagonals in wiggly pseudoquadrangles. We also orient this flip graph in a canonical way, which we call the *wiggly increasing flip graph* $WIFG_n$. The wiggly increasing flip graph on wiggly pseudotriangulations of 4 points is the front page illustration of this paper (throughout the paper, we implicitly orient the graphs and posets from bottom to top).

We then make a small detour to permutations. A *wiggly permutation* is a permutation of $[2n]$ avoiding the patterns $(2j-1) \cdots i \cdots (2j)$ for $i < 2j-1$ and $(2j) \cdots k \cdots (2j-1)$ for $k > 2j$. Our main result on wiggly permutations is the following.

Proposition. *The wiggly permutations induce a sublattice of the weak order on permutations of $[2n]$, which we call the *wiggly lattice* WL_n . Moreover, its Hasse diagram is regular of degree $2n-1$.*

We then connect wiggly pseudotriangulations to wiggly permutations.

Proposition. *The increasing flip graph on wiggly pseudotriangulations WIFG_n is isomorphic to the Hasse diagram of the wiggly lattice WL_n . In particular, the wiggly pseudotriangulations admit a lattice structure.*

Polyhedral geometry. We then switch to geometric realizations of the wiggly complex WC_n . Our constructions are inspired by similar constructions for generalized associahedra [HLT11, HPS18] of finite type cluster algebras [FZ02, FZ03], and for gentle associahedra [PPP21] of support τ -tilting complexes [AIR14] on gentle algebras [BR87]. Namely, we first construct a fan supported by certain \mathbf{g} -vectors associated to the wiggly arcs.

Theorem. *The cones generated by the \mathbf{g} -vectors of wiggly pseudodissections form a complete simplicial fan, called **wiggly fan** WF_n , realizing the wiggly complex WC_n .*

The proof of this result is based on a fine understanding of the linear dependences between \mathbf{g} -vectors. We further use this understanding to find a height function κ on wiggly arcs which satisfies all wall-crossing inequalities of the wiggly fan WF_n . Without entering the details of the definition of κ at the moment, let us just mention that it is obtained by summing some compatibility degrees of wiggly arcs, analogously to the constructions for cluster algebras [HPS18] and gentle algebras [PPP21]. Using the \mathbf{c} -vectors of each pseudotriangulation, defined to form the dual basis of the \mathbf{g} -vectors, we obtain our main result.

Theorem. *The wiggly fan WF_n is the normal fan of a simplicial $(2n - 1)$ -dimensional polytope, called the **wigglyhedron** W_n , and defined equivalently as*

- the intersection of the halfspaces $\{\mathbf{x} \in \mathbb{H} \mid \langle \mathbf{g}(\alpha) \mid \mathbf{x} \rangle \leq \kappa(\alpha)\}$ for all internal wiggly arcs α ,
- the convex hull of the points $\mathbf{p}(T) := \sum_{\alpha \in T} \kappa(\alpha) \mathbf{c}(\alpha, T)$ for all wiggly pseudotriangulations T .

Moreover, as for the classical permutahedron and associahedron, we recover our lattice structure by a suitable linear orientation of the graph of the wigglyhedron.

Proposition. *The Hasse diagram of the wiggly lattice WL_n (or equivalently, the wiggly increasing flip graph WIFG_n) is isomorphic to the graph of the wigglyhedron W_n oriented in a suitable direction.*

Further topics. The wigglyhedron has interesting connections in lattice theory and discrete geometry. We first connect the wigglyhedron to the (type A) Cambrian world.

Theorem. *For any $\delta \in \{-, +\}^n$,*

- the δ -Cambrian lattice of [Rea06] is an interval in the wiggly lattice WL_n ,
- the δ -Cambrian fan of [RS09] is a link in the wiggly fan WF_n ,
- the δ -associahedron of [HL07] is normally equivalent to a face of the wigglyhedron W_n .

We then define the wiggly complex WC_P of an arbitrary planar point set P (not necessarily aligned along the horizontal axis, nor necessarily in general position). Our definition recovers:

- when P consists of $n+2$ collinear points, the wiggly complex WC_n considered in this paper,
- when P is in general position, the pseudodissection complex of P considered in discrete and computational geometry [PV96, RSS08], and realized by the pseudotriangulation polytope in [RSS03] using rigidity and expansive motions.

These two families of examples motivate the following conjecture.

Conjecture. *For any point set P in the plane, the wiggly complex WC_P is the boundary complex of a simplicial polytope.*

We also present a few open questions related to this conjecture, concerning in particular the possible interpretation of the wiggly complex WC_P in terms of dual pseudoline arrangements, and the extension to multi wiggly complexes and arbitrary finite Coxeter groups.

Categorical interpretation. Each wiggly arc on $n + 2$ points corresponds to an object of an abelian category, such that morphisms and extensions between objects in the category are captured by minimal intersection numbers between the wiggly arcs. In fact, this correspondence holds for more general curves, which then correspond to objects of a larger triangulated category [KS02].

A general curve can be decomposed into wiggly arcs simply by cutting along its U-turns. We interpret this operation categorically as the decomposition of an object of the triangulated category into its cohomology pieces, from which we deduce the following.

Proposition. *The set of wiggly arcs corresponding to the cohomology pieces of an object represented by a general arc is a face of the wiggly complex.*

Furthermore, we characterize categorically when two objects in the abelian category are compatible, that is, their corresponding wiggly arcs are compatible.

Proposition. *Two wiggly arcs are compatible if and only if there are no extensions between the corresponding objects in the abelian category.*

2. WIGGLY PSEUDOTRIANGULATIONS AND WIGGLY PERMUTATIONS

Fix an integer $n \geq 1$. In this section, we define two combinatorial families, the wiggly pseudotriangulations of $n + 2$ points (Section 2.1) and the wiggly permutations of $[2n]$ (Section 2.2), and we prove that they are in bijection (Section 2.3). These two perspectives enable us to endow these combinatorial families with additional structures: the wiggly pseudotriangulations are actually the facets of the wiggly complex, while the wiggly permutations are actually the elements of the wiggly lattice. We conclude with a few questions and conjectures on graph properties of the wiggly flip graph (Section 2.4).

2.1. Wiggly pseudotriangulations and the wiggly complex. We start with wiggly pseudodissections of $n + 2$ points on a line. The following two definitions are illustrated in Figure 1.

Definition 1. A *wiggly arc* is a quadruple $\alpha := (i, j, A, B)$ where $0 \leq i < j \leq n + 1$ and the sets A and B form a partition of $]i, j[:= \{i + 1, \dots, j - 1\}$. We represent α by an x -monotone curve wiggling around the horizontal axis, starting at point i , ending at point j , and passing above the points of A and below the points of B . The wiggly arcs $\alpha_{\text{top}} := (0, n + 1, [n], \emptyset)$ and $\alpha_{\text{bot}} := (0, n + 1, \emptyset, [n])$ are called *external*, all other wiggly arcs are called *internal*.

Remark 2. The number of wiggly arcs is $\sum_{\ell=0}^n 2^\ell (n + 1 - \ell)$.

Definition 3. Two wiggly arcs (i, j, A, B) and (i', j', A', B') are

- *non pointed* if $i = j'$ or $i' = j$,
- *crossing* if $(A \cap B') \cup (\{i, j\} \cap B') \cup (A \cap \{i', j'\}) \neq \emptyset \neq (A' \cap B) \cup (\{i', j'\} \cap B) \cup (A' \cap \{i, j\})$, that is, if the corresponding curves cross,
- *incompatible* if they are non pointed or crossing, and *compatible* otherwise.

Remark 4. We want to stress out that our notion of compatibility on wiggly arcs of Definition 3 might seem similar to, but fundamentally differs from the notion of compatibility used by N. Reading to define noncrossing arc diagrams [Rea15]. Namely, our wiggly arcs and our crossing condition coincide with that of [Rea15]. However, we forbid an arc to start where another arc ends but allow two arcs to start or end at the same point, while N. Reading does precisely the opposite in [Rea15].

We then consider pairwise compatible sets of wiggly arcs. The following two definitions are illustrated in Figure 2.



FIGURE 1. Some incompatible wiggly arcs: non pointed (left) and crossing (right).




FIGURE 2. Three wiggly pseudodissections for $n = 7$. The left one has three wiggly cells, of respective degrees 3 (red), 8 (green), and 4 (blue). The other two have 7 pseudotriangles, and are obtained from each other by flipping the wiggly arc separating their yellow and dark blue pseudotriangles.

Definition 5. A *wiggly pseudodissection* D is a set of pairwise compatible wiggly arcs which contains the exterior wiggly arcs α_{top} and α_{bot} . We denote by $D^\circ := D \setminus \{\alpha_{\text{top}}, \alpha_{\text{bot}}\}$.

Definition 6. A wiggly pseudodissection D decomposes the digon bounded by α_{top} and α_{bot} into cells, that we call *wiggly cells*. The *degree* δ_c of a wiggly cell c is $\delta_c := \beta_c/2 + 2\kappa_c - 1$ where β_c is the number of wiggly arcs on the boundary of c (counted twice if they appear twice along the boundary of c), and κ_c is the number of connected components of the boundary of c . Two consecutive wiggly arcs along the boundary of c define a *corner* (resp. a *hinge*) of c if they bound a convex (resp. concave) angle of c .

Remark 7. Consider a wiggly cell c in a wiggly pseudodissection D . Note that β_c is always even as D is pointed, so that $\beta_c/2$ is an integer. Moreover, we have $\beta_c \geq 2$ and $\kappa_c \geq 1$, where at least one of the two inequalities is strict (as otherwise, the two wiggly arcs bounding c would coincide). We thus obtain on the one hand that $\delta_c = \beta_c/2 + 2\kappa_c - 1 \geq 3$ with equality if and only if $\beta_c = 4$ and $\kappa_c = 1$, and on the other hand that $\beta_c/2 + \kappa_c - 1 \geq 2$.

Definition 8. A *wiggly pseudotriangle* (resp. *pseudoquadrangle*) is a wiggly cell of degree 3 (resp. 4).

Remark 9. As illustrated in Figure 3, a wiggly pseudotriangle t has no point in its interior and 4 points on its boundary, 3 corners and 1 hinge. As illustrated in Figure 4, a wiggly pseudoquadrangle q can have

- (1) either one point in its interior and 2 points on its boundary, both corners,
- (2) or no point in its interior and 6 points on its boundary, 4 corners and 2 hinges. We then distinguish three situations, depending on the position of the two pairs p, p' of wiggly arcs incident to the two hinges of c , as illustrated in Figure 4 (where p and p' are colored in red and orange):
 - (a) those where p and p' are not consecutive along c , and do not overlap vertically,
 - (b) those where p and p' are not consecutive along c , but overlap vertically,
 - (c) those where p and p' are consecutive along c .

Proposition 10. Any wiggly pseudodissection D contains at most $2n - 1$ internal wiggly arcs and at most n wiggly cells. Moreover, the following are equivalent:

- (i) all wiggly cells of D are wiggly pseudotriangles,
- (ii) D contains n wiggly cells and is connected,
- (iii) D contains $2n - 1$ internal wiggly arcs,
- (iv) D is an inclusion maximal wiggly pseudodissection.

We then say that D is a *wiggly pseudotriangulation*.




FIGURE 3. Four wiggly pseudotriangles.




FIGURE 4. Four types of wiggly pseudoquadrangles described in Remark 9.

Proof. Consider a wiggly pseudodissection D with ι internal wiggly arcs, γ wiggly cells, and κ connected components. Euler's formula gives $(n+2) - (\iota+2) + (\gamma+1) = (\kappa+1)$, hence $n - \iota + \gamma = \kappa$. Moreover, $\sum_{c \in D} \beta_c/2 = 1 + \iota$ since each external (resp. internal) wiggly arc bounds precisely one (resp. two) wiggly cell of D , and $\sum_{c \in D} (\kappa_c - 1) = \kappa - 1$ since the boundary of each internal connected component of D is one of the internal connected components of the boundary of precisely one wiggly cell of D . As $2 \leq \beta_c/2 + \kappa_c - 1$ for any wiggly cell c by Remark 7, we thus obtain that $2\gamma \leq \sum_{c \in D} (\beta_c/2 + \kappa_c - 1) = \iota + \kappa = n + \gamma$. We conclude that $\gamma \leq n$ and $\iota = n + \gamma - \kappa \leq 2n - 1$.

To prove the second part of the statement, we show that (i) \Rightarrow (ii) \Rightarrow (iii) \Rightarrow (iv) \Rightarrow (i).

(i) \Rightarrow (ii). For any wiggly cell c , we have $\delta_c = 3$, so that $\beta_c = 4$ and $\kappa_c = 1$ by Remark 7, hence $2 = \beta_c/2 + \kappa_c - 1$. Therefore $\kappa = 1$ (D is connected) and $2\gamma = \iota + 1 = n + \gamma$, so that $\gamma = n$.

(ii) \Rightarrow (iii). As $\gamma = n$ and $\kappa = 1$, we have $\iota = n + \gamma - \kappa = 2n - 1$.


(iii) \Rightarrow (iv). If D already contains $2n - 1$ internal wiggly arcs, it has maximal cardinality among all wiggly pseudodissections, hence it is certainly inclusion maximal.

(iv) \Rightarrow (i). We now consider a wiggly pseudodissection D containing a wiggly cell c which is not a wiggly pseudotriangle, and we prove that D is not inclusion maximal. Assume first that the external boundary of c is a digon bounded by the wiggly arcs $\alpha := (i, j, A, B)$ and $\alpha' := (i, j, A', B')$. Denote by m the maximum of the points $(A \Delta A') \cup (B \Delta B')$ located inside this digon (where Δ denotes the symmetric difference). Then the wiggly arc $\beta := (i, m,]i, m[\cap (A \cup A'),]i, m[\cap (B \cap B'))$ is compatible with D (any wiggly arc incompatible with β crosses α or α') and is not in D (it connects the boundary of c with one of its internal connected components).

Assume now that the external boundary of c is a not digon. Consider a hinge j of c . Assume by symmetry that the boundary of c contains two wiggly arcs $\alpha := (i, j, A, B)$ and $\alpha' := (i', j, A', B')$, such that α is below α' . Let $\gamma := (k, \ell, X, Y)$ denote the uppermost wiggly arc of the boundary of c such that $j \in Y$ (it exists since j is a hinge of c), and let $X' :=]j, \ell[\cap X$ and $Y' :=]j, \ell[\cap Y$. Then the two wiggly arcs $\beta := (i, \ell, A \cup X', B \cup \{j\} \cup Y')$ and $\beta' := (i', \ell, A' \cup \{j\} \cup X', B \cup Y')$ are both compatible with D (any wiggly arc incompatible with β is incompatible with α or γ , and similarly for β') and cannot both belong to D (otherwise, c would be a wiggly pseudotriangle bounded by $\alpha, \alpha', \beta', \gamma$). \square

Remark 11. Note that in the proof of Proposition 10, we actually obtained that

$$\sum_{c \in C} (\delta_c - 3) = \sum_{c \in D} \beta_c / 2 + 2 \sum_{c \in D} (\kappa_c - 1) - 2\gamma = (1 + \iota) + 2(\kappa - 1) - 2(\iota + \kappa - n) = 2n - 1 - \iota.$$

Example 12. For $n = 1$, the 2 wiggly pseudotriangulations of 3 points are . For $n = 2$, the 14 wiggly pseudotriangulations of 4 points are



Example 13. The following wiggly pseudotriangulations are illustrated in Figure 5:

$$\begin{aligned} T_{\downarrow} &:= \{(i - 1, n + 1, \emptyset, [i, n]) \mid i \in [n + 1]\} \cup \{(0, n + 1, [i], [i, n]) \mid i \in [n]\}, \\ T_{\uparrow} &:= \{(0, i, \emptyset, [i - 1]) \mid i \in [n + 1]\} \cup \{(0, n + 1, [i, n], [i]) \mid i \in [n]\}, \\ T_{\leftarrow} &:= \{(0, 1, \emptyset, \emptyset)\} \cup \{(0, i + 1, \emptyset, [i]) \mid i \in [n]\} \cup \{(0, i + 1, [i], \emptyset) \mid i \in [n]\}, \\ T_{\rightarrow} &:= \{(n, n + 1, \emptyset, \emptyset)\} \cup \{(i - 1, n + 1, \emptyset, [i, n]) \mid i \in [n]\} \cup \{(i - 1, n + 1, [i, n], \emptyset) \mid i \in [n]\}. \end{aligned}$$




FIGURE 5. The wiggly pseudotriangulations T_{\downarrow} (left), T_{\uparrow} (middle left), T_{\leftarrow} (middle right), and T_{\rightarrow} (right) of Example 13 for $n = 7$.

Remark 14. The number wp_n of wiggly pseudotriangulations of $n + 2$ points is given by

n	1	2	3	4	5	6	7	8	...
wp_n	2	14	176	3232	78384	2366248	85534176	3602770400	...

To compute wp_n , denote by $wp_n(x)$ the polynomial where the coefficient of x^i is the number of wiggly pseudotriangulations of $n + 2$ points with i *final* internal wiggly arcs (*i.e.* ending at the last point $n + 1$). For instance,

$$\begin{aligned} wp_1(x) &= x + 1, \\ wp_2(x) &= 3x^3 + 5x^2 + 4x + 2, \\ wp_3(x) &= 15x^5 + 35x^4 + 44x^3 + 40x^2 + 28x + 14, \\ wp_4(x) &= 105x^7 + 315x^6 + 520x^5 + 630x^4 + 620x^3 + 514x^2 + 352x + 176. \end{aligned}$$

We invite the reader to check the expressions of $wp_1(x)$ and $wp_2(x)$ with Example 12. We use this additional variable x to obtain a recursive formula for $wp_n(x)$. Indeed, any wiggly pseudotriangulation of $n + 2$ points can be obtained from a wiggly pseudotriangulation of $n + 1$ points by inserting a single wiggly pseudotriangle with a hinge at n . Each wiggly pseudotriangulation of $n + 1$ points with i final internal wiggly arcs contributes to:

- $i + 2$ wiggly pseudotriangulations of $n + 2$ points with $i + 2$ final internal wiggly arcs, containing the wiggly arc $(n, n + 1, \emptyset, \emptyset)$,
- $j + 1$ wiggly pseudotriangulations of $n + 2$ points with j final internal wiggly arcs, not containing the wiggly arc $(n, n + 1, \emptyset, \emptyset)$, for all $0 \leq j \leq i + 1$.




FIGURE 6. The two diagonals in each of the four types of wiggly pseudoquadrangles.

This translates to the recursive formula

$$\begin{aligned}
 wp_n(x) &= x \cdot \frac{\partial}{\partial x} (x^2 wp_{n-1}(x)) + \frac{\partial}{\partial x} \left(\frac{wp_{n-1}(1) - x^3 wp_{n-1}(x)}{1-x} \right) \\
 &= \frac{1}{(1-x)^2} (wp_{n-1}(1) + x^2(2x^2 - 2x - 1)wp_{n-1}(x) + x^4(x-1)wp'_{n-1}(x)).
 \end{aligned}$$

This enables us to quickly compute $wp_n(x)$ and we obtain the numbers of wiggly pseudotriangulations wp_n by evaluating $wp_n(x)$ at $x = 1$.

Our next statement prepares for the definition of flips in wiggly pseudotriangulations, and is illustrated in Figure 6.

Definition 15. A *wiggly diagonal* of a wiggly cell c is a wiggly arc inside c and compatible with c .

Proposition 16. *Any wiggly pseudoquadrangle has exactly two wiggly diagonals. Moreover, these two wiggly diagonals either cross precisely once, or are non pointed.*

Proof. Depending on the four types of pseudoquadrangles described in Remark 9:

- (1) the two diagonals are the two wiggly arcs inside q , incident to the point in the interior of q ,
- (2a) the two diagonals connect the two pairs of opposite corners of q ,
- (2b) one of two diagonals connects two opposite corners of q while the other connects the two hinges of q ,
- (2c) one of the diagonals connects two opposite corners of q while the other connects an hinge of q to a corner of q .

See Figure 6 where the two diagonals appear in red and orange. \square

We are now ready to consider the wiggly flip graph and the wiggly complex. The following definition is illustrated in Figure 7 (right) when $n = 2$.

Definition 17. The *wiggly flip graph* WFG_n is the graph with a vertex for each wiggly pseudo-triangulation, and with an edge between two wiggly pseudotriangulations T and T' if there are wiggly arcs $\alpha \in T$ and $\alpha' \in T'$ such that $T \setminus \{\alpha\} = T' \setminus \{\alpha'\}$.

Proposition 18. *The wiggly flip graph WFG_n is regular of degree $2n - 1$ and connected.*

Proof. Consider any internal wiggly arc α of a wiggly pseudotriangulation T . Then $Q := T \setminus \{\alpha\}$ contains $n - 2$ wiggly pseudotriangles and 1 pseudoquadrangle q . By Proposition 16, q admits precisely two wiggly diagonals, namely α and another one α' . As α' is inside q and compatible with c , it is also compatible with Q . We obtain that $T' := Q \cup \{\alpha'\}$ is the only wiggly pseudo-triangulation other than T containing Q . This shows that any internal wiggly arc in any wiggly pseudotriangulation can be flipped, hence that the wiggly flip graph is regular of degree $2n - 1$ by Proposition 10 (iii).

There are various possible proofs for the connectedness. For instance, it will follow easily from Proposition 38. To avoid relying on this forward reference, we sketch an alternative direct proof




FIGURE 7. The wiggly complex WC_2 (left) and the wiggly flip graph WFG_2 (right).

by induction on n . We claim that if a wiggly pseudotriangulation has a non-zero number of final internal wiggly arcs (*i.e.* ending at the last point $n+1$), then at least one of them can be flipped to a non-final wiggly arc. This implies that any wiggly pseudotriangulation can be transformed by flips to a wiggly pseudotriangulation with no final internal wiggly arc. All wiggly pseudotriangulations with no final internal wiggly arc contain the arcs $(0, n, \emptyset, [n-1])$ and $(0, n, [n-1], \emptyset)$, since the latter are compatible with all internal wiggly arcs but the final ones. Hence, they induce a subgraph of WFG_n isomorphic to WFG_{n-1} , which is connected by induction. We conclude that WFG_n is itself connected.



To see the claim, let i be maximal such that T contains two distinct wiggly arcs starting at i and ending at $n+1$ (it is well-defined as we have the external wiggly arcs α_{top} and α_{bot} for $i=0$). Consider a final internal wiggly arc α of T starting weakly after i , and let α' be the wiggly arc obtained by flipping α in T . If α' is final, then by Proposition 16, at least one pseudotriangle t of T incident to α has two corners at $n+1$. Let h be the hinge of t and β, β' be the two wiggly arcs of t joining h to $n+1$. As $i < h \leq n$ the maximality of i imposes that $\beta = \beta'$. Flipping $\beta = \beta'$ in T thus yields a wiggly arc ending at $h \leq n$, hence a non-final arc. \square

We will also consider an oriented version of the wiggly flip graph WFG_n . The following definition is illustrated in Figure 8.

Definition 19. Let $\alpha := (i, j, A, B)$ and $\alpha' := (i', j', A', B')$ be two wiggly arcs and T and T' be two wiggly pseudotriangulations, such that $T \setminus \{\alpha\} = T' \setminus \{\alpha'\}$. We say that the flip from T to T' is *increasing* if

- α starts where α' ends (that is, $i = j'$), or
- α crosses α' from north-west to south-east (that is, there exists $k \in [n-1]$ such that $k \in (A \cap B') \cup (\{i\} \cap B') \cup (A \cap \{i'\})$ and $k+1 \in (A' \cap B) \cup (\{j'\} \cap B) \cup (A' \cap \{j\})$).

The *wiggly increasing flip graph* $WIFG_n$ is the directed graph with a vertex for each wiggly pseudotriangulation and an edge for each increasing flip.

FIGURE 8. The wiggly arcs α and α' of Definition 19.FIGURE 9. The wiggly arcs β and β' of Proposition 20.

We will need the following observation about exchangeable arcs, illustrated in Figure 9.

Proposition 20. *Let $\alpha := (i, j, A, B)$ and $\alpha' := (i', j', A', B')$ be two exchangeable wiggly arcs. Then any wiggly arc compatible with both α and α' is also compatible with the wiggly arcs:*

- $\beta := (i, j', A \cup A', B \cup B' \cup \{j\})$ and $\beta' := (i, j', A \cup A' \cup \{j\}, B \cup B')$ if α and α' are non pointed with $j = i'$, (In other words, β and β' are the two wiggly arcs starting at $\min(i, i')$ and ending at $\max(j, j')$ which follow α on $]i, j[$ and α' on $]i', j'[,$)
- $\beta := (i, j',]i, j'[\setminus (B \cup B'),]i, j'[\cap (B \cup B'))$ and $\beta' := (i', j,]i', j[\cap (A \cup A'),]i', j[\setminus (A \cup A'))$ if α and α' are crossing and there is $k \in [n-1]$ such that $k \in (A \cap B') \cup (\{i\} \cap B') \cup (A \cap \{i')$ and $k+1 \in (A' \cap B) \cup (\{j'\} \cap B) \cup (A' \cap \{j\})$. (In other words, β is the wiggly arc starting at i and ending at j' which follows the lower hull of α and α' , and β' is the wiggly arc starting at i' and ending at j which follows the upper hull of α and α' .)

(By exchanging α and α' , similar statements hold if α and α' are non pointed with $j = i'$, or if α and α' are crossing with $i \in B'$ or $i' \in A$.)

Proof. Any wiggly arc non pointed with β would be non pointed with α or α' (as the endpoints of β are endpoints of α or α'). Any wiggly arc crossing β would be either non pointed or crossing α or α' (as β follows α or α' , except at the point $j = i'$ in the first situation). \square

Finally, we define the wiggly complex WC_n , illustrated in Figure 7 (left) when $n = 2$.

Definition 21. The *wiggly complex* WC_n is the simplicial complex of pairwise pointed and non-crossing subsets of internal wiggly arcs. In other words, it is the clique complex of the compatibility graph on internal wiggly arcs.

Proposition 22. *The wiggly complex WC_n is a flag $(2n - 1)$ -dimensional pseudomanifold without boundary.*

Proof. It is a flag simplicial complex because it is a clique complex. It is pure of dimension $2n - 1$ by the implication (iv) \Rightarrow (iii) of Proposition 10. Finally, it is a pseudomanifold without boundary by Proposition 18. \square

Remark 23. WC_1 contains two isolated points. WC_2 is a simplicial 3-dimensional associahedron (this coincidence fails for $n > 2$). Here are the first few f -vectors:

$$\begin{aligned} f(WC_1) &= (1, 2), \\ f(WC_2) &= (1, 9, 21, 14), \\ f(WC_3) &= (1, 24, 154, 396, 440, 176), \\ f(WC_4) &= (1, 55, 729, 4002, 10930, 15684, 11312, 3232), \\ f(WC_5) &= (1, 118, 2868, 28110, 140782, 400374, 673274, 662668, 352728, 78384), \end{aligned}$$

and the first few h -vectors:

$$\begin{aligned} h(WC_1) &= (1, 1), \\ h(WC_2) &= (1, 6, 6, 1), \\ h(WC_3) &= (1, 19, 68, 68, 19, 1), \\ h(WC_4) &= (1, 48, 420, 1147, 1147, 420, 48, 1), \\ h(WC_5) &= (1, 109, 1960, 11254, 25868, 25868, 11254, 1960, 109, 1). \end{aligned}$$

Note that the number of vertices and facets of WC_n were already discussed in Remarks 2 and 14.

Remark 24. The vertical and horizontal reflections provide two obvious automorphisms of the wiggly complex WC_n (and thus of the wiggly flip graph WFG_n). Moreover, the vertical reflection is an anti-automorphism of the wiggly increasing flip graph $WIFG_n$ (while the horizontal reflection is not).

2.2. Wiggly permutations and the wiggly lattice. We now consider wiggly permutations of $[2n]$, defined as follows.

Definition 25. A *wiggly permutation* is a permutation of $[2n]$ which avoids the patterns:

- $(2j-1) \cdots i \cdots (2j)$ for $j \in [n]$ and $i < 2j-1$,
- $(2j) \cdots k \cdots (2j-1)$ for $j \in [n]$ and $k > 2j$.

Example 26. For $n = 1$, the 2 wiggly permutations of $[2]$ are 12 and 21. For $n = 2$, the 14 wiggly permutations of $[4]$ are

$$1234, 1243, 1342, 1423, 1432, 2134, 2143, 3412, 3421, 4123, 4132, 4213, 4312, 4321.$$

Example 27. The permutations

$$\sigma_{\downarrow} := 12 \cdots (2n), \quad \sigma_{\uparrow} := (2n) \cdots 21, \quad \sigma_{\leftarrow} := (2n) \cdots 21 \cdots (2n-1), \quad \sigma_{\rightarrow} := 1 \cdots (2n-1)(2n) \cdots 2$$

are wiggly permutations of $[2n]$.

Remark 28. The number wp_n of wiggly permutations of $[2n]$ is given by

n	1	2	3	4	5	6	7	8	...
wp_n	2	14	176	3232	78384	2366248	85534176	3602770400	...

To compute wp_n , denote by $wp_n(x)$ the polynomial where the coefficient of x^i is the number of wiggly permutations of $[2n]$ with i *admissible gaps* (i.e. gaps γ between two consecutive positions such that there is no $j \in [n]$ such that the value $2j$ appears before the gap γ while the value $2j-1$ appears after the gap γ). For instance,

$$\begin{aligned} wp_1(x) &= x + 1, \\ wp_2(x) &= 3x^3 + 5x^2 + 4x + 2, \\ wp_3(x) &= 15x^5 + 35x^4 + 44x^3 + 40x^2 + 28x + 14, \\ wp_4(x) &= 105x^7 + 315x^6 + 520x^5 + 630x^4 + 620x^3 + 514x^2 + 352x + 176. \end{aligned}$$

We invite the reader to check the expressions of $wp_1(x)$ and $wp_2(x)$ with Example 26. Any wiggly permutation of $[2n]$ can be obtained from a wiggly permutation of $[2n-2]$ by inserting the last values $2n-1$ and $2n$. Each wiggly permutation of $[2n-2]$ with i admissible gaps contributes to:




FIGURE 10. The wiggly lattice WL_2 on wiggly pseudotriangulations (left) and on wiggly permutations (right).

- $i + 2$ wiggly permutations of $[2n]$ with $i + 2$ admissible gaps and where $2n - 1$ appears before $2n$ (they must appear consecutively),
- $j + 1$ wiggly permutations of $[2n]$ with j admissible arcs and where $2n$ appears before $2n - 1$, for all $0 \leq j \leq i + 1$.

We thus obtain the same recursive equation on $wp_n(x)$ as in Remark 14.

Recall that the *inversion set* and *non-inversion set* of a permutation σ of $[2n]$ are the sets

$$\begin{aligned} \text{inv}(\sigma) &:= \{(\sigma(i), \sigma(j)) \mid 1 \leq i < j \leq 2n \text{ and } \sigma(i) > \sigma(j)\}, \\ \text{ninv}(\sigma) &:= \{(\sigma(i), \sigma(j)) \mid 1 \leq i < j \leq 2n \text{ and } \sigma(i) < \sigma(j)\}. \end{aligned}$$

Note that $\text{inv}(\sigma)$ and $\text{ninv}(\sigma)$ are transitive: $\{(c, b), (b, a)\} \subseteq \text{inv}(\sigma)$ implies $(c, a) \in \text{inv}(\sigma)$ for all $1 \leq a < b < c \leq 2n$, and similarly for ninv . The inversion sets of wiggly permutations are characterized as follows.

Lemma 29. *A permutation σ of $[2n]$ is a wiggly permutation if and only if for all $j \in [n]$,*

- $(2j - 1, i) \in \text{inv}(\sigma)$ implies $(2j, i) \in \text{inv}(\sigma)$ for all $i < 2j - 1$, and
- $(k, 2j - 1) \in \text{inv}(\sigma)$ implies $(k, 2j) \in \text{inv}(\sigma)$ for all $k > 2j$.

A similar characterization holds for $\text{ninv}(\sigma)$ by exchanging $2j - 1$ and $2j$.

Proof. Immediate as it corresponds to the pattern avoidance description of Definition 25. \square

The (left) *weak order* on permutations of $[2n]$ is defined as the inclusion order of their inversion sets (or equivalently, the reverse inclusion of their non-inversion sets). Its cover relations are given by the exchanges of two entries at consecutive positions. It is a (congruence uniform) lattice, and the join and meet satisfy

$$\text{inv}(\sigma \vee \tau) = (\text{inv}(\sigma) \cup \text{inv}(\tau))^{\text{tc}} \quad \text{and} \quad \text{ninv}(\sigma \wedge \tau) = (\text{ninv}(\sigma) \cup \text{ninv}(\tau))^{\text{tc}}$$

where X^{tc} denotes the transitive closure of X .

The following statements are illustrated on Figure 10 (right) when $n = 2$.

Proposition 30. *The wiggly permutations induce a sublattice of the weak order on permutations of $[2n]$, that we call the *wiggly lattice* WL_n .*

Proof. Consider two wiggly permutations ρ and σ of $[2n]$, and let $\tau = \rho \vee \sigma$. Assume that there is $1 \leq i < 2j - 1 < 2n$ such that $(2j - 1, i) \in \text{inv}(\tau)$. As $\text{inv}(\tau) = (\text{inv}(\rho) \cup \text{inv}(\sigma))^{\text{tc}}$, there exist $i \leq i' < 2j - 1$ such that $(i', i) \in \text{inv}(\tau)$ and $(2j - 1, i') \in \text{inv}(\rho) \cup \text{inv}(\sigma)$. Since ρ and σ are wiggly permutations, we obtain by Lemma 29 that $(2j, i') \in \text{inv}(\rho) \cup \text{inv}(\sigma) \subseteq \text{inv}(\tau)$. As $(i', i) \in \text{inv}(\tau)$ and $(2j, i') \in \text{inv}(\tau)$ and $\text{inv}(\tau)$ is transitive, we conclude that $(2j, i) \in \text{inv}(\tau)$. A similar argument shows that $(k, 2j - 1) \in \text{inv}(\tau)$ implies $(k, 2j) \in \text{inv}(\tau)$ for all $1 < 2j < k \leq 2n$. By Lemma 29, we conclude that $\tau = \rho \vee \sigma$ is a wiggly permutation. The proof is similar for $\rho \wedge \sigma$, using nin instead of inv . \square

Remark 31. The map $\sigma_1 \dots \sigma_{2n} \mapsto (2n + 1 - \sigma_1) \dots (2n + 1 - \sigma_n)$ obviously provides an anti-isomorphism of the wiggly lattice WL_n .

We now aim at describing the cover relations of the wiggly lattice. Recall that an *ascent* (resp. *descent*) in a permutation σ is a position j such that $\sigma(j) < \sigma(j + 1)$ (resp. $\sigma(j) > \sigma(j + 1)$).

Lemma 32. *Consider an ascent j of a wiggly permutation σ of $[2n]$. Let $i := \min(j, \sigma^{-1}(\sigma(j) + 1))$ if $\sigma(j)$ is odd, and $i := j$ otherwise. Let $k := \max(j + 1, \sigma^{-1}(\sigma(j + 1) + 1))$ if $\sigma(j + 1)$ is odd, and $k := j + 1$ otherwise. Then*

$$\sigma^j := \sigma(1) \dots \sigma(i - 1) \sigma(j + 1) \dots \sigma(k) \sigma(i) \dots \sigma(j) \sigma(k + 1) \dots \sigma(2n).$$

is the minimal wiggly permutation such that $\text{inv}(\sigma) \cup \{(\sigma(j + 1), \sigma(j))\} \subseteq \text{inv}(\sigma^j)$.

Proof. We start with a few observations. Let $i < p \leq j$ (resp. $j + 1 \leq q < k$). As σ is a wiggly permutation, we have $\sigma(p) < \sigma(i)$ (resp. $\sigma(q) > \sigma(k)$). Hence $(\sigma(i), \sigma(p))$ (resp. $(\sigma(q), \sigma(k))$) belongs to $\text{inv}(\sigma)$. Moreover, since j is an ascent of σ , we obtain that $\sigma(p) < \sigma(q)$ for all $i \leq p \leq j < q \leq k$. Hence, $\text{inv}(\sigma^j) = \text{inv}(\sigma) \cup \{(\sigma(q), \sigma(p)) \mid i \leq p \leq j < q \leq k\}$.

We now prove that σ^j is indeed a wiggly permutation of $[2n]$. We consider $v \in [n]$ and $u < 2v - 1$ such that $(2v - 1, u) \in \text{inv}(\sigma^j)$, and we show that $(2v, u) \in \text{inv}(\sigma^j)$. If $(2v, u) \in \text{inv}(\sigma)$, we are done as $\text{inv}(\sigma) \subset \text{inv}(\sigma^j)$. If $(2v, u) \notin \text{inv}(\sigma)$, then $(2v - 1, u) \notin \text{inv}(\sigma)$ by Lemma 29, and $i \leq \sigma^{-1}(u) \leq j < \sigma^{-1}(2v - 1) \leq k$ since $\text{inv}(\sigma^j) \setminus \text{inv}(\sigma) = \{(\sigma(q), \sigma(p)) \mid i \leq p \leq j < q \leq k\}$. We have $\sigma^{-1}(2v) \leq k$ since σ is a wiggly permutation. As $(2v, u) \notin \text{inv}(\sigma)$ and $\sigma(p) < \sigma(q)$ for all $i \leq p \leq j < q \leq k$, we conclude that $j < \sigma^{-1}(2v) < k$. Hence, we obtain that $i \leq \sigma^{-1}(u) \leq j < \sigma^{-1}(2v) \leq k$, so that $(2v, u) \in \text{inv}(\sigma^j)$. Similarly $(w, 2v - 1) \in \text{inv}(\sigma^j)$ implies $(w, 2v) \in \text{inv}(\sigma^j)$ for all $v \in [n]$ and $w > 2v$. We conclude that σ^j is indeed a wiggly permutation of $[2n]$ by Lemma 29.

Consider now any wiggly transposition τ such that $\text{inv}(\sigma) \cup \{(\sigma(j + 1), \sigma(j))\} \subseteq \text{inv}(\tau)$. By Lemma 29, we obtain that $\text{inv}(\tau)$ also contains $(\sigma(k), \sigma(i))$. Since $\text{inv}(\sigma) \subset \text{inv}(\tau)$ and $\text{inv}(\sigma)$ contains $(\sigma(i), \sigma(p))$ and $(\sigma(q), \sigma(k))$ for all $i < p \leq j$ and $j + 1 \leq q < k$, we conclude by transitivity that $\text{inv}(\tau)$ contains $(\sigma(q), \sigma(p))$ for all $i \leq p \leq j < q \leq k$. \square

Proposition 33. *Each wiggly permutation σ covers (resp. is covered by) as many wiggly permutations as its number of descents (resp. ascents). Hence, the cover graph of WL_n is regular of degree $2n - 1$ and connected.*

Proof. Observe that $\sigma^j \neq \sigma^{j'}$ for distinct ascents $j \neq j'$. As any permutation σ' larger than σ satisfies $\text{inv}(\sigma') \supseteq \text{inv}(\sigma) \cup \{(\sigma(j + 1), \sigma(j))\}$ for some ascent j of σ , we thus obtain that $j \rightarrow \sigma^j$ is a bijection from the ascents of σ to the wiggly permutations covering σ . The proof is symmetrical for the bijection from descents of σ to wiggly permutations covered by σ . \square

2.3. Bijection. We now prove that wiggly pseudotriangulations and wiggly permutations are in bijection, and that this bijection induces an isomorphism from the wiggly flip graph to the cover graph of the wiggly lattice. The following two definitions are illustrated in Figures 10 and 11.

Definition 34. Consider a wiggly pseudotriangulation T . Rotating counterclockwise inside each wiggly pseudotriangle of T , we label by $2h - 1$ (resp. $2h + 1$) the corner immediately preceding (resp. following) the hinge h (the remaining corner remains unlabelled). We define $\Phi(T)$ as the permutation obtained by reading these labels from bottom to top, meaning that for each internal wiggly arc α of T , the label incident to α and below α appears just before the label incident to α and above α .




FIGURE 11. The bijection between wiggly pseudotriangulations and wiggly permutations.

Example 35. The map Φ sends the wiggly pseudotriangulations of Example 13 to the wiggly permutations of Example 27. Namely, $\Phi(T_{\downarrow}) = \sigma_{\downarrow}$, $\Phi(T_{\uparrow}) = \sigma_{\uparrow}$, $\Phi(T_{\leftarrow}) = \sigma_{\leftarrow}$, and $\Phi(T_{\rightarrow}) = \sigma_{\rightarrow}$.

Definition 36. For a wiggly permutation σ of $[2n]$, we define $\Psi(\sigma) := \{\alpha(\sigma, k) \mid k \in [2n-1]\}$, where for $k \in [2n-1]$, we have $\alpha(\sigma, k) := (i, j, A, B)$ with

$$\begin{aligned} i &:= \max\{0\} \cup \{i \in [n] \mid 2i-1 \in \sigma([k]) \not\preceq 2i\}, \\ j &:= \min\{n+1\} \cup \{j \in [n] \mid 2j \in \sigma([k]) \not\preceq 2j-1\}, \\ A &:= \{\ell \in]i, j[\mid \{2\ell-1, 2\ell\} \subseteq \sigma([k])\}, \\ B &:= \{\ell \in]i, j[\mid \{2\ell-1, 2\ell\} \cap \sigma([k]) = \emptyset\}. \end{aligned}$$

Example 37. The map Ψ sends the wiggly permutations of Example 27 to the wiggly pseudotriangulations of Example 13. Namely, $\Psi(\sigma_{\downarrow}) = T_{\downarrow}$, $\Psi(\sigma_{\uparrow}) = T_{\uparrow}$, $\Psi(\sigma_{\leftarrow}) = T_{\leftarrow}$, and $\Psi(\sigma_{\rightarrow}) = T_{\rightarrow}$.

Proposition 38. The maps Φ of Definition 34 and Ψ of Definition 36 define inverse bijections between the wiggly pseudotriangulations and the wiggly permutations, which induce a directed graph isomorphism between the wiggly increasing flip graph WIFG_n and the Hasse diagram of the wiggly lattice WL_n .

Proof. First, we show that $\Phi(T)$ is a well-defined wiggly permutation for each wiggly pseudotriangulation T . Observe first that $\Phi(T)$ is by construction a permutation σ of $[2n]$. For any $j \in [n]$, let t_j be the pseudotriangle of T that has hinge j . Suppose that $2j-1$ appears before $2j$ in σ for some fixed $j \in [n]$. This means that the pseudotriangle t_j has its unlabelled corner to the left of its labelled corners. In other words, its hinge opens to the right (see *e.g.* t_2 in Figure 11). Consider any label i that we encounter as we traverse from $2j-1$ to $2j$. This label belongs to some pseudotriangle t_h with $h > j$. In other words, σ avoids the pattern $2j-1 \cdots i \cdots 2j$ for $i < 2j-1$. The other case (in which $2j$ appears before $2j-1$ in σ) is symmetric.

Next, we show that $\Psi(\sigma)$ is a wiggly pseudotriangulation for each wiggly permutation σ . Observe first that for any $1 \leq u < v \leq n$, we cannot see $\{2u, 2v-1\}$ before $\{2u-1, 2v\}$ in σ as it contains no forbidden pattern of Definition 25. Using the notations of Definition 36, this implies that $i < j$ and that $]i, j[= A \sqcup B$, so that $\alpha(\sigma, k) := (i, j, A, B)$ is indeed a wiggly arc for each $k \in [2n-1]$. By Proposition 10, it thus suffices to check that for each $k \neq k'$ in $[2n-1]$, the wiggly arcs $\alpha(\sigma, k)$ and $\alpha(\sigma, k')$ are distinct and compatible.

- (1) **Distinctness.** Suppose by contradiction that $\alpha(\sigma, k) = (i, j, A, B) = \alpha(\sigma, k')$ for some $k < k'$. Let $\ell \in \sigma([k'] \setminus [k])$ and $\bar{\ell} := \lceil \ell/2 \rceil$. As $\ell \notin \sigma([k])$, we have $\bar{\ell} \notin A$. As $\ell \in \sigma([k'])$, we have $\bar{\ell} \notin B$. As $\{2i-1, 2j\} \subseteq \sigma([k])$ and $\{2i, 2j-1\} \cap \sigma([k']) = \emptyset$ and σ avoids the forbidden pattern of Definition 25, we have $2i < \ell < 2j-1$, so that $i < \bar{\ell} < j$. This contradicts the fact that $]i, j[= A \sqcup B$.
- (2) **Compatibility.** Let $\alpha(\sigma, k) = (i, j, A, B)$ and $\alpha(\sigma, k') = (i', j', A', B')$ for some $k < k'$. It is clear from the definition that $i \neq j'$, as the sets over which we take the max and min to define i and j' respectively are disjoint. Similarly, $j \neq i'$. Hence, the two wiggly arcs are pointed. To check that the wiggly arcs are non-crossing, we show that $(A \cap B') \cup$

$(\{i, j\} \cap B') \cup (A \cap \{i', j'\}) = \emptyset$. Indeed $A \cap B' = \emptyset$ since $A \subseteq A'$ because $\sigma([k]) \subseteq \sigma([k'])$. Moreover, $\{i, j\} \cap B' = \emptyset$ since $\{2i - 1, 2j\} \subseteq \sigma([k]) \subseteq \sigma([k'])$. Finally, $A \cap \{i', j'\} = \emptyset$ since $\{2i', 2j' - 1\} \cap \sigma([k]) \subseteq \{2i', 2j' - 1\} \cap \sigma([k']) = \emptyset$.

Next, we show that the maps Φ and Ψ are inverse bijections. Let σ be a wiggly permutation, so that $\Psi(\sigma)$ is a wiggly pseudotriangulation. Let $k \in [2n - 1]$ and $m := \lceil \sigma(k)/2 \rceil$. Then observe that

- if $\{2m, 2m - 1\} \subseteq \sigma([k])$, then k is minimal such that $\alpha(\sigma, k)$ passes above the point m ,
- if $\sigma(k) = 2m$ and $2m - 1 \notin \sigma([k])$, then k is minimal such that $\alpha(\sigma, k)$ has right endpoint m ,
- if $\sigma(k) = 2m - 1$ and $2m \notin \sigma([k])$, then k is minimal such that $\alpha(\sigma, k)$ has left endpoint m .

It follows that in the wiggly pseudotriangulation $\Psi(\sigma)$, the wiggly arc $\alpha(\sigma, k)$ is precisely the wiggly arc lying immediately above the angle labelled by $\sigma(k)$. Therefore, we have $\Phi(\Psi(\sigma)) = \sigma$.

Conversely, let T be a wiggly pseudotriangulation, so that $\sigma = \Phi(T)$ is a wiggly permutation. Then for any $k \in [2n - 1]$, the wiggly arc immediately above the angle of T labelled by $\sigma(k)$ is precisely the one constructed as $\alpha(\sigma, k)$. This shows that $\Psi(\Phi(T)) = T$.

Finally, consider a cover relation $\sigma < \sigma^j$ in the wiggly lattice WL_n as described in Lemma 32. Using the notations of Lemma 32, observe that

- $\alpha(\sigma^j, \ell) = \alpha(\sigma, \ell)$ for all $\ell < i$ or $\ell > k$,
- $\alpha(\sigma^j, \ell) = \alpha(\sigma, \ell + k - j)$ for all $i \leq \ell < j$, and
- $\alpha(\sigma^j, \ell) = \alpha(\sigma, \ell - j + i - 1)$ for all $j < \ell \leq k$.

Hence, $\Psi(\sigma) \setminus \{\alpha(\sigma, j)\} = \Psi(\sigma^j) \setminus \{\alpha(\sigma^j, j)\}$. This implies that Ψ is a directed graph morphism from the Hasse diagram of the wiggly lattice WL_n to the wiggly increasing flip graph WIFG_n . As the underlying undirected graphs are both regular of degree $2n - 1$, we conclude that Φ and Ψ induce directed graph isomorphisms between the wiggly increasing flip graph WIFG_n and the Hasse diagram of the wiggly lattice WL_n . \square

Remark 39. Transporting the wiggly lattice WL_n via the map Ψ to the wiggly pseudotriangulation, note that the minimal wiggly pseudotriangulation is $\Psi(\sigma_\downarrow) = T_\downarrow$ while the maximal wiggly pseudotriangulation is $\Psi(\sigma_\uparrow) = T_\uparrow$. See Example 13 and Figure 5 for illustrations.

Remark 40. The map Φ sends the wiggly complex automorphism of Remark 24 given by the vertical reflection to the wiggly lattice anti-isomorphism of Remark 31.

Remark 41. The map Φ sends wiggly pseudotriangulations with i final internal wiggly arcs to wiggly permutations with i admissible gaps, explaining the similarities of Remarks 14 and 28.

2.4. Graph properties. We conclude this combinatorial section with two open questions and a conjecture on graphical properties of the wiggly flip graph WFG_n (or equivalently of the cover graph of the wiggly lattice WL_n), motivated by the relevance of similar questions for the associahedron.

First, we are interested in the graph diameter δ_n of WFG_n . It is clearly bounded below by $2n - 1$ (as $T_\leftarrow \cap T_\rightarrow = \{\alpha_{\text{top}}, \alpha_{\text{bot}}\}$, all $2n - 1$ internal wiggly arcs must sometimes be flipped at least once) and above by $\binom{2n}{2}$ (as we have a sublattice of the weak order on permutations of $[2n]$). The first few values of δ_n are

n	1	2	3	4	5	...
δ_n	1	4	8	14	20	...

but they might be misleading as for the associahedron [Pou14]. This raises the following question.

Question 42. Compute (or estimate) the diameter δ_n of the wiggly flip graph WFG_n .

Related to the estimation of the diameter δ_n is the study of the properties of the geodesic paths in WFG_n . We refer to [CP16] for a discussion on geodesic properties in a polytope, in connection to the revisiting path property and the complexity of the simplex algorithm [San13, San12].

Question 43. Does the wiggly complex has the non-leaving face property of [CP16]?

Finally, we checked the following conjecture computationally up to $n = 3$.

Conjecture 44. *The wiggly flip graph WFG_n admits an Hamiltonian path (or even cycle).*

Note that unfortunately, wiggly permutations do not form a zigzag language in the sense of [HHMW22] (which would have guaranteed the existence of a Gray code).

3. WIGGLY FAN AND WIGGLYHEDRON

In this section, we use \mathbf{g} - and \mathbf{c} -vectors (Section 3.2) to define the wiggly fan (Section 3.3) and the wigglyhedron (Section 3.4), which provide geometric realizations of the wiggly complex and of the wiggly lattice.

3.1. Polyhedral geometry. We refer to [Zie98] for a reference on polyhedral geometry, and only remind the basic notions needed later in the paper.

A (polyhedral) **cone** is the positive span $\mathbb{R}_{\geq 0}\mathbf{R}$ of a finite set \mathbf{R} of vectors of \mathbb{R}^d or equivalently, the intersection of finitely many closed linear half-spaces of \mathbb{R}^d . The **faces** of a cone are its intersections with its supporting hyperplanes. The **rays** (resp. **facets**) are the faces of dimension 1 (resp. codimension 1). A cone is **simplicial** if its rays are linearly independent. A (polyhedral) **fan** \mathcal{F}_D is a set of cones such that any face of a cone of \mathcal{F}_D belongs to \mathcal{F}_D , and any two cones of \mathcal{F}_D intersect along a face of both. A fan is **essential** if the intersection of its cones is the origin, **complete** if the union of its cones covers \mathbb{R}^d , and **simplicial** if all its cones are simplicial.

Note that a simplicial fan defines a simplicial complex on its rays (the simplices of the simplicial complex are the subsets of rays which span a cone of the fan). Conversely, given a simplicial complex Δ with ground set V , one can try to realize it geometrically by associating a ray \mathbf{r}_v of \mathbb{R}^d to each $v \in V$, and the cone $\mathbb{R}_{\geq 0}\mathbf{R}_\Delta$ generated by the set $\mathbf{R}_\Delta := \{\mathbf{r}_v \mid v \in \Delta\}$ to each $\Delta \in \Delta$. To show that the resulting cones indeed form a fan, we will need the following statement, which can be seen as a reformulation of [DRS10, Coro. 4.5.20].

Proposition 45. *Consider a simplicial $(d-1)$ -dimensional pseudomanifold Δ without boundary on a ground set V and a set of vectors $(\mathbf{r}_v)_{v \in V}$ of \mathbb{R}^d , and define $\mathbf{R}_\Delta := \{\mathbf{r}_v \mid v \in \Delta\}$ for any $\Delta \in \Delta$. Then the collection of cones $\{\mathbb{R}_{\geq 0}\mathbf{R}_\Delta \mid \Delta \in \Delta\}$ forms a complete simplicial fan of \mathbb{R}^d if and only if*

- *there exists a vector \mathbf{v} of \mathbb{R}^d contained in only one of the cones $\mathbb{R}_{\geq 0}\mathbf{R}_\Delta$ for $\Delta \in \Delta$,*
- *for any two adjacent facets Δ, Δ' of Δ with $\Delta \setminus \{v\} = \Delta' \setminus \{v'\}$, we have $\lambda_v \lambda_{v'} > 0$ where*

$$\lambda_v \mathbf{r}_v + \lambda_{v'} \mathbf{r}_{v'} + \sum_{w \in \Delta \cap \Delta'} \lambda_w \mathbf{r}_w = 0$$

denotes the unique (up to rescaling) linear dependence on $\mathbf{R}_{\Delta \cup \Delta'}$.

A **polytope** is the convex hull of finitely many points of \mathbb{R}^d or equivalently, a bounded intersection of finitely many closed affine half-spaces of \mathbb{R}^d . The **faces** of a polytope are its intersections with its supporting hyperplanes. The **vertices** (resp. **edges**, resp. **facets**) are the faces of dimension 0 (resp. dimension 1, resp. codimension 1).

The **normal cone** of a face \mathbb{F} of a polytope \mathbb{P} is the cone generated by the normal vectors to the supporting hyperplanes of \mathbb{P} containing \mathbb{F} . Said differently, it is the cone of vectors \mathbf{c} of \mathbb{R}^d such that the linear form $\mathbf{x} \mapsto \langle \mathbf{c} \mid \mathbf{x} \rangle$ on \mathbb{P} is maximized by all points of the face \mathbb{F} . The **normal fan** of \mathbb{P} is the set of normal cones of all its faces.

Consider now a complete simplicial fan \mathcal{F}_D of \mathbb{R}^d with rays $(\mathbf{r}_v)_{v \in V}$ and cones $\mathbb{R}_{\geq 0}\mathbf{R}_\Delta$ for $\Delta \in \Delta$, where $\mathbf{R}_\Delta := \{\mathbf{r}_v \mid v \in \Delta\}$ as in Proposition 45. To realize the fan \mathcal{F}_D , one can try to pick a height vector $\mathbf{h} := (h_v)_{v \in V} \in \mathbb{R}^V$ and consider the polytope $\mathbb{P}_\mathbf{h} := \{\mathbf{x} \in \mathbb{R}^d \mid \langle \mathbf{r}_v \mid \mathbf{x} \rangle \leq h_v \text{ for all } v \in V\}$. The following classical statement characterizes the height vectors \mathbf{h} for which the fan \mathcal{F}_D is the normal fan of this polytope $\mathbb{P}_\mathbf{h}$. We borrow the formulation from [CFZ02, Lem. 2.1].

Proposition 46. *Let \mathcal{F}_D be an essential complete simplicial fan in \mathbb{R}^n with rays $(\mathbf{r}_v)_{v \in V}$ and cones $\mathbb{R}_{\geq 0}\mathbf{R}_\Delta$ for $\Delta \in \Delta$. Then the following are equivalent for any height vector $\mathbf{h} \in \mathbb{R}^V$:*

- *The fan \mathcal{F}_D is the normal fan of the polytope $\mathbb{P}_\mathbf{h} := \{\mathbf{x} \in \mathbb{R}^d \mid \langle \mathbf{r}_v \mid \mathbf{x} \rangle \leq h_v \text{ for all } v \in V\}$.*
- *For two adjacent facets Δ, Δ' of Δ with $\Delta \setminus \{v\} = \Delta' \setminus \{v'\}$, the height vector \mathbf{h} satisfies the wall crossing inequality*

$$\lambda_v h_v + \lambda_{v'} h_{v'} + \sum_{w \in \Delta \cap \Delta'} \lambda_w h_w > 0,$$

where

$$\lambda_v \mathbf{r}_v + \lambda_{v'} \mathbf{r}_{v'} + \sum_{w \in \Delta \cap \Delta'} \lambda_w \mathbf{r}_w = 0$$

denotes the unique linear dependence on $\mathbf{R}_{\Delta \cup \Delta'}$ such that $\lambda_v + \lambda_{v'} = 2$.




FIGURE 12. The \mathbf{g} -matrix $\mathbf{g}(T)$ and 4 times the \mathbf{c} -matrix $\mathbf{c}(T)$ of a wiggly pseudotriangulation T . The i th column of these matrices are the \mathbf{g} -vector $\mathbf{g}(\alpha)$ and 4 times the \mathbf{c} -vector $\mathbf{c}(\alpha, T)$ of the i th wiggly arc α of T (ordered from bottom to top). The colors of the columns match the colors of the wiggly arcs.

We denote by $(\mathbf{e}_i)_{i \in [d]}$ the standard basis of \mathbb{R}^d . For $I \subseteq [d]$, we define $\mathbf{1}_I := \sum_{i \in I} \mathbf{e}_i$, and we often shorten $\mathbf{1}_{[d]}$ by $\mathbf{1}_d$. We denote by \mathbb{H}_d the hyperplane of \mathbb{R}^d defined by the equation $\langle \mathbf{x} \mid \mathbf{1}_d \rangle = 0$, and we denote by $\pi : \mathbb{R}^d \rightarrow \mathbb{H}$ the orthogonal projection on \mathbb{H}_d , that is $\pi(\mathbf{x}) := \mathbf{x} - (\langle \mathbf{x} \mid \mathbf{1}_d \rangle / d) \mathbf{1}_d$.

3.2. \mathbf{g} -vectors and \mathbf{c} -vectors. We now define two families of vectors, illustrated in Figure 12.

Definition 47. The \mathbf{g} -vector $\mathbf{g}(\alpha)$ of a wiggly arc $\alpha := (i, j, A, B)$ is defined as the projection $\pi(\hat{\mathbf{g}}(\alpha))$ of the vector $\hat{\mathbf{g}}(\alpha) := \mathbf{1}_{\alpha^+} - \mathbf{1}_{\alpha^-}$ where

$$\begin{aligned} \alpha^+ &:= (\{2i - 1, 2j\} \setminus \{-1, 2n + 2\}) \cup \{2a - 1 \mid a \in A\} \cup \{2a \mid a \in A\}, \\ \alpha^- &:= (\{2i, 2j - 1\} \setminus \{0, 2n + 1\}) \cup \{2b - 1 \mid b \in B\} \cup \{2b \mid b \in B\}. \end{aligned}$$

If we place the coordinate $2p - 1$ on the left and the coordinate $2p$ on the right of point p , then $\mathbf{g}(\alpha)$ has a 1 outside its two endpoints and on both sides of points in A , and a -1 inside its two endpoints and on both sides of points in B . Note that $\hat{\mathbf{g}}((0, n + 1, [n], \emptyset)) = \mathbf{1}_{2n} = -\hat{\mathbf{g}}((0, n + 1, \emptyset, [n]))$, so that $\mathbf{g}((0, n + 1, [n], \emptyset)) = \mathbf{0} = \mathbf{g}((0, n + 1, \emptyset, [n]))$. We define $\mathbf{g}(D) := \{\mathbf{g}(\alpha) \mid \alpha \in D^\circ\}$ for a set D of wiggly arcs.

Example 48. Table 1 gathers the \mathbf{g} -matrices (*i.e.* matrices whose columns are the \mathbf{g} -vectors) of the wiggly pseudotriangulations of Example 13 illustrated in Figure 5.

Observe that the \mathbf{g} -matrices of Example 48 have full rank. More generally, we will see in Corollary 56 that $\mathbf{G}(T)$ forms a basis of \mathbb{H}_{2n} for any wiggly pseudotriangulation T , which enables us to define its dual basis.

Definition 49. The \mathbf{c} -vector of an interior wiggly arc α in a wiggly pseudotriangulation T is the vector $\mathbf{c}(\alpha, T)$ of \mathbb{H}_{2n} such that $\langle \mathbf{c}(\alpha, T) \mid \mathbf{g}(\alpha) \rangle = 1$ and $\langle \mathbf{c}(\alpha, T) \mid \mathbf{g}(\alpha') \rangle = 0$ for all interior wiggly arcs $\alpha' \neq \alpha$ of T . In other words, $\mathbf{c}(T) := \{\mathbf{c}(\alpha, T) \mid \alpha \in T^\circ\}$ is the dual basis of $\mathbf{g}(T)$ in \mathbb{H}_{2n} .

Remark 50. The \mathbf{c} -vector $\mathbf{c}(\alpha, T)$ can also be described combinatorially. Namely, label the the corners of T as in Definition 34, and denote by u and v the labels incident to α and respectively below and above α , and let $\bar{u} := \lceil u/2 \rceil$ and $\bar{v} := \lceil v/2 \rceil$. Then the \mathbf{c} -vector $\mathbf{c}(\alpha, T)$ has coordinates

- $\mathbf{c}(\alpha, T)_{2w-1} = -\mathbf{c}(\alpha, T)_{2w} = (-1)^{w \in A} (-1)^{u > v} / 4$ for all w strictly between \bar{u} and \bar{v} ,
- $\mathbf{c}(\alpha, T)_u = 1/2$ if $\bar{u} = \bar{v}$, or if \bar{v} and the wiggly arcs incident to \bar{u} are on the same side of \bar{u} , and $\mathbf{c}(\alpha, T)_{2\bar{u}-1} = \mathbf{c}(\alpha, T)_{2\bar{u}} = 1/4$ otherwise,
- $\mathbf{c}(\alpha, T)_v = -1/2$ if $\bar{u} = \bar{v}$, or if \bar{u} and the wiggly arcs incident to \bar{v} are on the same side of \bar{v} , and $\mathbf{c}(\alpha, T)_{2\bar{v}-1} = \mathbf{c}(\alpha, T)_{2\bar{v}} = -1/4$ otherwise,
- $\mathbf{c}(\alpha, T)_w = 0$ for all other coordinates.

The tedious proof of this description is not needed here as we only use the dual basis property.

Example 51. Table 2 gathers the \mathbf{c} -matrices (*i.e.* matrices whose columns are the \mathbf{c} -vectors) of the wiggly pseudotriangulations of Example 13 illustrated in Figure 5.

Remark 52. Following Remark 24, note that the vertical and horizontal symmetries on wiggly arcs translate on \mathbf{g} - and \mathbf{c} -vectors to the isometries sending $(x_1, x_2, \dots, x_{2n-1}, x_{2n}) \in \mathbb{H}_{2n}$ to $(x_{2n}, x_{2n-1}, \dots, x_2, x_1)$ and to $(-x_2, -x_1, \dots, -x_{2n}, -x_{2n-1})$ respectively.

3.3. Wiggly fan. We now show that the \mathbf{g} -vectors of Definition 47 support a polyhedral fan realization of the wiggly complex WC_n .

Theorem 53. *The collection of cones $\mathbb{R}_{\geq 0}\mathbf{g}(D)$ for all wiggly dissections D in the wiggly complex WC_n forms a complete simplicial fan of \mathbb{H}_{2n} , called the **wiggly fan** WF_n .*

To prove Theorem 53, we just need to check the conditions of Proposition 45. We first find a vector contained in the cone $\mathbb{R}_{\geq 0}\mathbf{g}(T)$ for a single wiggly pseudotriangulation T .

Lemma 54. *Let $\mathbf{v} := \sum_{k \in [n]} (\mathbf{e}_{2k} - \mathbf{e}_{2k-1}) = (-1, 1, \dots, -1, 1)$, let $\alpha_0 := (0, 1, \emptyset, \emptyset)$, and for $i \in [n]$, let $\underline{\alpha}_i := (0, i+1, [i], \emptyset)$ and $\bar{\alpha}_i := (0, i+1, \emptyset, [i])$. Then $T_{\leftarrow} := \{\alpha_0\} \cup \bigcup_{i \in [n]} \{\underline{\alpha}_i, \bar{\alpha}_i\}$ is the only wiggly pseudotriangulation T such that $\mathbf{v} \in \mathbb{R}_{\geq 0}\mathbf{g}(T)$.*

Proof. Observe first that, for any wiggly arc $\alpha := (i, j, A, B)$ and any $k \in [n]$, we have

$$\hat{\mathbf{g}}(\alpha)_{2k-1} - \hat{\mathbf{g}}(\alpha)_{2k} = \begin{cases} 1 & \text{if } k = i, \\ -1 & \text{if } k = j, \\ 0 & \text{otherwise,} \end{cases} \quad \text{and} \quad \hat{\mathbf{g}}(\alpha)_{2k-1} + \hat{\mathbf{g}}(\alpha)_{2k} = \begin{cases} 2 & \text{if } k \in A, \\ -2 & \text{if } k \in B, \\ 0 & \text{otherwise.} \end{cases}$$

Assume now that $\mathbf{v} = \sum_{\alpha} \lambda_{\alpha} \hat{\mathbf{g}}(\alpha)$ where the sum ranges over all wiggly arcs α for some positive scalars $\lambda_{\alpha} \geq 0$ such that $\lambda_{\alpha} \cdot \lambda_{\alpha'} = 0$ for all incompatible wiggly arcs α and α' . For $k \in [n]$, let $\underline{\Lambda}_k$ (resp. $\bar{\Lambda}_k$) denote the sum of λ_{α} over the wiggly arcs $\alpha := (i, j, A, B)$ with $k \in A$ (resp. $k \in B$). Note that $\underline{\Lambda}_k = \bar{\Lambda}_k$ since $0 = \mathbf{v}_{2k-1} + \mathbf{v}_{2k} = \sum_{\alpha} \lambda_{\alpha} (\hat{\mathbf{g}}(\alpha)_{2k-1} + \hat{\mathbf{g}}(\alpha)_{2k}) = 2\underline{\Lambda}_k - 2\bar{\Lambda}_k$.

As $\mathbf{v}_1 - \mathbf{v}_2 = -2$ and $\hat{\mathbf{g}}(\alpha)_1 - \hat{\mathbf{g}}(\alpha)_2 \geq 0$ except if $\alpha = \alpha_0$, we have $\lambda_{\alpha_0} > 0$, and thus $\lambda_{\alpha} = 0$ for all wiggly arcs α starting at 1. We then prove by induction on $i \in [n-1]$ that $\lambda_{\underline{\alpha}_i} \cdot \lambda_{\bar{\alpha}_i} > 0$ and that $\lambda_{\alpha} = 0$ for all wiggly arcs starting at $i+1$. Consider $i \in [n-1]$ and assume that we proved it for $i-1$. As $\lambda_{\underline{\alpha}_{i-1}} \cdot \lambda_{\bar{\alpha}_{i-1}} > 0$, we already have $\lambda_{\alpha} = 0$ for all wiggly arcs α ending at $i+1$, except if $\alpha \in \{\underline{\alpha}_i, \bar{\alpha}_i\}$. As $\mathbf{v}_{2i+1} - \mathbf{v}_{2i+2} = -2$, this implies that $\lambda_{\underline{\alpha}_i} + \lambda_{\bar{\alpha}_i} = 1$. This already implies that $\lambda_{\alpha} = 0$ for all wiggly arcs α starting at $i+1$. Moreover, assume for instance that $\lambda_{\bar{\alpha}_i} > 0$. Hence, $\lambda_{\alpha} = 0$ for all wiggly arcs $\alpha := (i, j, A, B)$ with $i \in A$ and $i+1 \in B$. We thus obtain that $\underline{\Lambda}_i - \lambda_{\underline{\alpha}_i} \leq \underline{\Lambda}_{i+1}$ and $\bar{\Lambda}_i - \lambda_{\bar{\alpha}_i} \geq \bar{\Lambda}_{i+1}$. As $\underline{\Lambda}_i = \bar{\Lambda}_i$ and $\underline{\Lambda}_{i+1} = \bar{\Lambda}_{i+1}$, we thus obtain that $\lambda_{\underline{\alpha}_i} \geq \lambda_{\bar{\alpha}_i} > 0$, hence that $\lambda_{\underline{\alpha}_i} \cdot \lambda_{\bar{\alpha}_i} > 0$. We conclude that $\lambda_{\alpha} = 0$ for all wiggly arcs α except if $\alpha = \alpha_0$ or $\alpha \in \{\underline{\alpha}_i, \bar{\alpha}_i\}$ for some $i \in [n]$.

Consider now a wiggly pseudotriangulation T such that $\mathbf{v} \in \mathbb{R}_{\geq 0}\mathbf{g}(T)$. Write $\mathbf{v} = \sum_{\alpha \in T^{\circ}} \lambda_{\alpha} \mathbf{g}(\alpha)$ with $\lambda_{\alpha} \geq 0$ for all $\alpha \in T^{\circ}$. Let $\mu := \sum_{\alpha \in T^{\circ}} \lambda_{\alpha} \langle \hat{\mathbf{g}}(\alpha) | \mathbf{1}_{2n} \rangle$. As $\mathbf{v} = \sum_{\alpha \in T^{\circ}} \lambda_{\alpha} \hat{\mathbf{g}}(\alpha) - \mu \mathbf{1}_{2n}$ and $\mathbf{1}_{2n} = \hat{\mathbf{g}}((0, n+1, [n], \emptyset)) = -\hat{\mathbf{g}}((0, n+1, \emptyset, [n]))$, we thus conclude from the previous paragraph that $T = T_{\leftarrow}$. \square

We now describe the unique linear dependence between the \mathbf{g} -vectors of two adjacent facets of the wiggly complex WC_n . The following statement is illustrated in Figure 9.

Lemma 55. *Let T and T' be two adjacent wiggly pseudotriangulations with $T \setminus \{\alpha\} = T' \setminus \{\alpha'\}$, and let β and β' be the wiggly arcs defined in Proposition 20. Then $\{\beta, \beta'\} \in T \cap T'$ and*

- if α and α' are not pointed, then $\mathbf{g}(\alpha) + \mathbf{g}(\alpha') = (\mathbf{g}(\beta) + \mathbf{g}(\beta'))/2$,
- if α and α' are crossing, then $\mathbf{g}(\alpha) + \mathbf{g}(\alpha') = \mathbf{g}(\beta) + \mathbf{g}(\beta')$.

Proof. In both situations, we will just prove that the same linear dependence holds on the vectors $\hat{\mathbf{g}}(\alpha), \hat{\mathbf{g}}(\alpha'), \hat{\mathbf{g}}(\beta), \hat{\mathbf{g}}(\beta')$, which will imply the statement as π is a linear map.

If α and α' are non pointed, then we have

$$\alpha^+ \sqcup \alpha'^+ = \beta^+ \sqcup \{2j-1, 2j\} = \beta'^+ \quad \text{and} \quad \alpha^- \sqcup \alpha'^- = \beta^- = \beta'^- \sqcup \{2j-1, 2j\}.$$

We thus obtain that

$$\mathbf{1}_{\alpha^+} + \mathbf{1}_{\alpha'^+} = \mathbf{1}_{\beta^+} + \mathbf{e}_{2j-1} + \mathbf{e}_{2j} = \mathbf{1}_{\beta'^+} \quad \text{and} \quad \mathbf{1}_{\alpha^-} + \mathbf{1}_{\alpha'^-} = \mathbf{1}_{\beta^-} = \mathbf{1}_{\beta'^-} + \mathbf{e}_{2j-1} + \mathbf{e}_{2j}.$$

hence

$$\begin{aligned} 2(\hat{\mathbf{g}}(\alpha) + \hat{\mathbf{g}}(\alpha')) &= 2(\mathbf{1}_{\alpha^+} - \mathbf{1}_{\alpha^-}) + 2(\mathbf{1}_{\alpha'^+} - \mathbf{1}_{\alpha'^-}) = 2(\mathbf{1}_{\alpha^+} + \mathbf{1}_{\alpha'^+}) - 2(\mathbf{1}_{\alpha^-} + \mathbf{1}_{\alpha'^-}) \\ &= (\mathbf{1}_{\beta^+} + \mathbf{1}_{\beta'^+} + \mathbf{e}_{2j-1} + \mathbf{e}_{2j}) - (\mathbf{1}_{\beta^-} + \mathbf{1}_{\beta'^-} + \mathbf{e}_{2j-1} + \mathbf{e}_{2j}) \\ &= (\mathbf{1}_{\beta^+} + \mathbf{1}_{\beta'^+}) - (\mathbf{1}_{\beta^-} + \mathbf{1}_{\beta'^-}) = (\mathbf{1}_{\beta^+} - \mathbf{1}_{\beta^-}) + (\mathbf{1}_{\beta'^+} - \mathbf{1}_{\beta'^-}) = \hat{\mathbf{g}}(\beta) + \hat{\mathbf{g}}(\beta'). \end{aligned}$$

If α and α' cross, then we have

$$\begin{aligned} \alpha^+ \cup \alpha'^+ &= \beta^+ \cup \beta'^+ & \text{and} & \quad \alpha^+ \cap \alpha'^+ = \beta^+ \cap \beta'^+, \\ \alpha^- \cup \alpha'^- &= \beta^- \cup \beta'^- & \text{and} & \quad \alpha^- \cap \alpha'^- = \beta^- \cap \beta'^-. \end{aligned}$$

We thus obtain that

$$\mathbf{1}_{\alpha^+} + \mathbf{1}_{\alpha'^+} = \mathbf{1}_{\alpha^+ \cup \alpha'^+} + \mathbf{1}_{\alpha^+ \cap \alpha'^+} = \mathbf{1}_{\beta^+ \cup \beta'^+} + \mathbf{1}_{\beta^+ \cap \beta'^+} = \mathbf{1}_{\beta^+} + \mathbf{1}_{\beta'^+}$$

and similarly $\mathbf{1}_{\alpha^-} + \mathbf{1}_{\alpha'^-} = \mathbf{1}_{\beta^-} + \mathbf{1}_{\beta'^-}$, so that $\hat{\mathbf{g}}(\alpha) + \hat{\mathbf{g}}(\alpha') = \hat{\mathbf{g}}(\beta) + \hat{\mathbf{g}}(\beta')$. \square

Corollary 56. *For any wiggly pseudotriangulation T , the set $\mathbf{g}(T) := \{\mathbf{g}(\alpha) \mid \alpha \in T^\circ\}$ is a basis of \mathbb{H}_{2n} .*

Proof. Observe that the \mathbf{g} -matrix $\mathbf{g}(T_{\leftarrow})$ presented in Example 48 has full rank (it is very easy to check for $\mathbf{g}(T_{\leftarrow})$, but any other would do). The statement thus follows from Lemma 55 and the connectedness of the flip graph on wiggly pseudotriangulations. \square

Finally, we wrap up Theorem 53.

Proof of Theorem 53. Directly follows from Proposition 45 and Lemmas 54 and 55. \square

Remark 57. Following Remarks 24 and 52, note that the vertical and horizontal symmetries on wiggly arcs define linear automorphisms of the wiggly fan WF_n .

Remark 58. By definition, each wall of the wiggly fan is normal to a \mathbf{c} -vector. Hence, the wiggly fan coarsens the \mathbf{c} -vector fan, defined by the arrangement of hyperplanes $\{\mathbf{x} \in \mathbb{H}_{2n} \mid \langle \mathbf{x} \mid \mathbf{c}(\alpha, T) \rangle\}$ for all wiggly arcs α and wiggly pseudotriangulations T with $\alpha \in T$.

3.4. Wigglyhedron. We now construct the wigglyhedron illustrated in Figure 13 when $n = 2$.

Definition 59. The *incompatibility degree* $\delta(\alpha, \alpha')$ of two wiggly arcs $\alpha := (i, j, A, B)$ and $\alpha' := (i', j', A', B')$ is given by

- 0 if α and α' are pointed and non-crossing,
- 1 if α and α' are not pointed (*i.e.* $i = j'$ or $i' = j$),
- the number of crossings of α and α' if they are crossing (*i.e.* the number of $p \in \{0, \dots, n\}$ with $p \in (A \cap B') \cup (\{i, j\} \cap B') \cup (A \cap \{i', j'\})$ and $p+1 \in (A' \cap B) \cup (\{i', j'\} \cap B) \cup (A' \cap \{i, j\})$, or the opposite).

The *incompatibility number* of a wiggly arc α is $\kappa(\alpha) := \sum_{\alpha'} \delta(\alpha, \alpha')$.

Remark 60. Note that δ is indeed an incompatibility degree in the sense that


- α and α' are compatible if and only if $\delta(\alpha, \alpha') = 0$,
- α and α' are exchangeable if and only if $\delta(\alpha, \alpha') = 1$.

Observe also that δ (and thus κ) is preserved by the automorphisms of Remark 24.

Example 61. Observe that $\kappa((0, j, [1, j], \emptyset)) = \kappa((0, j, \emptyset, [1, j])) = (2^j - 1)(2^{n+1-j} - 1)$.

Lemma 62. *Let T and T' be two adjacent wiggly pseudotriangulations with $T \setminus \{\alpha\} = T' \setminus \{\alpha'\}$, and let β and β' be the wiggly arcs defined in Proposition 20. Then*

- if α and α' are not pointed, then $\kappa(\alpha) + \kappa(\alpha') > (\kappa(\beta) + \kappa(\beta'))/2$,
- if α and α' are crossing, then $\kappa(\alpha) + \kappa(\alpha') > \kappa(\beta) + \kappa(\beta')$.

FIGURE 13. The wigglyhedron W_2 .

Proof. Any crossing between a wiggly arc γ and β or β' translates to a crossing with α or α' . When α and α' are not pointed, a crossing with α or α' can correspond to a crossing with β and a crossing with β' , so that $\delta(\alpha, \gamma) + \delta(\alpha', \gamma) \geq (\delta(\beta, \gamma) + \delta(\beta', \gamma))/2$. When α and α' are crossing, then there is a correspondence between the crossings with α or α' and the crossings with β or β' , so that $\delta(\alpha, \gamma) + \delta(\alpha', \gamma) \geq \delta(\beta, \gamma) + \delta(\beta', \gamma)$. Moreover, both inequalities are strict for $\gamma = \alpha'$ since $\delta(\alpha, \alpha') = 1$ while $\delta(\beta, \alpha') = \delta(\beta', \alpha') = 0$. The result thus follows by summation. \square

Theorem 63. *The wiggly fan WF_n is the normal fan of a simplicial $(2n - 1)$ -dimensional polytope, called the **wigglyhedron** W_n , and defined equivalently as*

- the intersection of the halfspaces $\{\mathbf{x} \in \mathbb{H} \mid \langle \mathbf{g}(\alpha) \mid \mathbf{x} \rangle \leq \kappa(\alpha)\}$ for all internal wiggly arcs α ,
- the convex hull of the points $\mathbf{p}(T) := \sum_{\alpha \in T} \kappa(\alpha) \mathbf{c}(\alpha, T)$ for all wiggly pseudotriangulations T .

Proof. By Proposition 46 and Lemmas 55 and 62, the wiggly fan is indeed the normal fan of the polytope W_n defined by the inequalities $\langle \mathbf{g}(\alpha) \mid \mathbf{x} \rangle \leq \kappa(\alpha)$ for all internal wiggly arcs α . For any wiggly pseudotriangulation T and any wiggly arc α of T , we have $\langle \mathbf{g}(\alpha) \mid \mathbf{p}(T) \rangle = \sum_{\alpha' \in T} \kappa(\beta) \langle \mathbf{g}(\alpha) \mid \mathbf{c}(\alpha', T) \rangle = \kappa(\alpha)$ since $\mathbf{g}(T)$ and $\mathbf{c}(T)$ are dual bases by Definition 49. Hence, $\mathbf{p}(T)$ is indeed the point of W_n corresponding to T , which is located at the intersection of the hyperplanes $\langle \mathbf{g}(\alpha) \mid \mathbf{x} \rangle = \kappa(\alpha)$ for all internal wiggly arcs α in T . \square

In fact, the wigglyhedron even recovers the wiggly lattice of Section 2.2.

Proposition 64. *The Hasse diagram of the wiggly lattice WL_n is isomorphic to the graph of the wigglyhedron W_n oriented in the direction $\boldsymbol{\omega} := \sum_{i \in [n]} 4ni(\mathbf{e}_{2i-1} + \mathbf{e}_{2i}) + \sum_{j \in [2n]} j\mathbf{e}_j$.*

Proof. Consider an increasing flip $T \rightarrow T'$ between two wiggly pseudotriangulations T and T' with $T \setminus \{\alpha\} = T' \setminus \{\alpha'\}$. Denote by u and v the labels incident to α and respectively below and above α in the corner labeling of T described in Definition 34, and let $\bar{u} := \lceil u/2 \rceil$ and $\bar{v} := \lceil v/2 \rceil$. We have $u < v$ because the flip $T \rightarrow T'$ is increasing, so that $\bar{u} \leq \bar{v}$.

Observe now that $\mathbf{p}(T') - \mathbf{p}(T)$ is orthogonal to the $(2n - 2)$ -dimensional cone of the wiggly fan generated by the \mathbf{g} -vectors of the wiggly arcs in $T \cap T'$, and points from the side containing $\mathbf{g}(\alpha)$

to the side containing $\mathbf{g}(\alpha')$. Hence, $\mathbf{p}(T') - \mathbf{p}(T)$ is a positive multiple of $-\mathbf{c}(\alpha, T)$. From Definition 49, we observe that

- $\langle \mathbf{c}(\alpha, T) \mid \mathbf{e}_{2i-1} + \mathbf{e}_{2i} \rangle = 1/2$ if $i = \bar{u} \neq \bar{v}$, $-1/2$ if $i = \bar{v} \neq \bar{u}$, and 0 otherwise,
- $|\langle \mathbf{c}(\alpha, T) \mid \mathbf{e}_j \rangle| \leq 1/2$ for all $j \in [2n]$.

We now distinguish two situations. If $\bar{u} \neq \bar{v}$, then

$$\langle -\mathbf{c}(\alpha, T) \mid \boldsymbol{\omega} \rangle = \sum_{i \in [n]} 4ni \langle -\mathbf{c}(\alpha, T) \mid \mathbf{e}_{2i-1} + \mathbf{e}_{2i} \rangle + \sum_{j \in [2n]} j \langle -\mathbf{c}(\alpha, T) \mid \mathbf{e}_j \rangle \geq 2n(\bar{v} - \bar{u}) - n > 0.$$

If $\bar{u} = \bar{v}$, then $u = 2k - 1$ and $v = 2k$ for some $k \in [n]$, and we have

$$\langle \mathbf{c}(\alpha, T) \mid \boldsymbol{\omega} \rangle = \sum_{i \in [n]} 4ni \langle \mathbf{c}(\alpha, T) \mid \mathbf{e}_{2i-1} + \mathbf{e}_{2i} \rangle + \sum_{j \in [2n]} j \langle \mathbf{c}(\alpha, T) \mid \mathbf{e}_j \rangle = \frac{2k}{2} - \frac{2k-1}{2} = 1/2 > 0.$$

We conclude in both cases that $\langle \mathbf{c}(\alpha, T) \mid \boldsymbol{\omega} \rangle > 0$, so that we recover the increasing flip graph by orienting the graph of the wigglyhedron \mathbb{W}_n in direction $\boldsymbol{\omega}$. \square

Remark 65. Following Remarks 24, 52 and 57, note that the vertical and horizontal symmetries on wiggly arcs define isometries of the wigglyhedron \mathbb{W}_n .

4. FURTHER TOPICS

In this section, we explore two further topics on the wigglyhedron. We first observe that any (type A) Cambrian associahedron [Rea06, HL07] is normally equivalent to a face of the wigglyhedron \mathbb{W}_n (Section 4.1). We then raise the question to extend our construction to wiggly complexes of arbitrary planar point sets (Section 4.2).

4.1. Cambrian considerations. In this section, we fix a signature $\delta := \delta_1 \dots \delta_n \in \{+, -\}^n$. By convention, we set $\delta_0 = \delta_{n+1} = 0$. We connect the δ -Cambrian lattice [Rea06] to the wiggly lattice of Section 2 and the δ -associahedron of [HL07] to the wigglyhedron of Section 3. Definitions 66 to 69 and Proposition 72 are illustrated in Figure 14.

Definition 66. The δ -gon is a convex $(n+2)$ -gon with a point with abscissa i and with ordinate of the same sign as δ_i for all $0 \leq i \leq n+1$. A δ -triangulation is a triangulation of the δ -gon. The δ -triangulation lattice is the transitive closure of the slope increasing flip graph on δ -triangulations.

Definition 67. A δ -permutation is a permutation of $[n]$ which avoids $\dots ik \dots j \dots$ with $i < j < k$ and $\delta_j = -$, and $\dots j \dots ki \dots$ with $i < j < k$ and $\delta_j = +$. The δ -permutation lattice is the sublattice of the weak order induced by δ -permutations.

Definition 68. The δ -wiggly arcs are the arcs $(0, j, [1, j], \emptyset)$ for $\delta_j = +$ and $(0, j, \emptyset, [1, j])$ for $\delta_j = -$. A δ -wiggly pseudotriangulation is a wiggly pseudotriangulation containing all δ -wiggly arcs. The δ -wiggly pseudotriangulation lattice is the transitive closure of the increasing flip graph on δ -wiggly pseudotriangulations.

Definition 69. A δ -wiggly permutation is a wiggly permutation σ of $[2n]$ such that

$$\delta_j = + \implies \sigma^{-1}(i) \leq \sigma^{-1}(2j-1) \quad \text{and} \quad \delta_j = - \implies \sigma^{-1}(2j) \leq \sigma^{-1}(i)$$

for all $1 \leq i \leq 2j \leq 2n$. The δ -wiggly permutation lattice is the interval of the wiggly permutation lattice WL_n given by δ -wiggly permutations.

Remark 70. We had to make a choice in Definitions 68 and 69. For $\delta_j = +$, we could replace “ $(0, j, [1, j], \emptyset)$ ” by “ $(j, n+1, [j, n], \emptyset)$ ” in Definition 68, and “ $\sigma^{-1}(i) \leq \sigma^{-1}(2j-1)$ for $1 \leq i \leq 2j$ ” by “ $\sigma^{-1}(i) \leq \sigma^{-1}(2j-1)$ for $2j-1 \leq i \leq 2n$ ” in Definition 69. Independently, for $\delta_j = -$, we could replace “ $(0, j, \emptyset, [1, j])$ ” by “ $(j, n+1, \emptyset, [j, n])$ ” in Definition 68, and “ $\sigma^{-1}(2j) \leq \sigma^{-1}(i)$ for $1 \leq i \leq 2j$ ” by “ $\sigma^{-1}(2j-1) \leq \sigma^{-1}(i)$ for $2j-1 \leq i \leq 2n$ ” in Definition 69. Propositions 72 and 73 would still hold.

Remark 71. Following Remarks 24, 52, 57 and 65, note that the vertical and horizontal symmetries send $\delta := \delta_1 \dots \delta_n$ to $\delta_n \dots \delta_1$ and $(-\delta_1) \dots (-\delta_n)$ respectively.




FIGURE 14. Cambrian lattices as intervals of wiggly lattices.

Proposition 72. *For any signature $\delta \in \{+, -\}^n$, the δ -triangulation lattice, δ -permutation lattice, δ -wiggly pseudotriangulation lattice, and δ -wiggly permutation lattice are all isomorphic, and are known as the δ -Cambrian lattice [Rea06].*

Proof. We only describe the following bijections

$$\begin{array}{ccc}
 \delta\text{-triangulations} & \longleftarrow & \delta\text{-permutations} \\
 \downarrow & & \uparrow \\
 \delta\text{-wiggly pseudotriangulations} & \longrightarrow & \delta\text{-wiggly permutations}
 \end{array}$$

illustrated in Figure 14. We let the readers convince themselves that they are well-defined, that they compose to the identity, and that they preserve the flip structure. We also invite the reader to work out the explicit description of the 8 missing arrows. For $I \subseteq [n]$, we set $I^+ := \{i \in I \mid \delta_i = +\}$ and $I^- := \{i \in I \mid \delta_i = -\}$.

From δ -triangulations to δ -wiggly pseudotriangulations. Replace each diagonal (i, j) of the δ -gon by the wiggly arc $(0, j, [i] \cup]i, j[^- ,]i, j[^+)$ if $\delta_i \geq 0$ and $(0, j,]i, j[^-, [i] \cup]i, j[^+)$ if $\delta_i \leq 0$.

From δ -wiggly pseudotriangulations to δ -wiggly permutations. Specialization of Definition 34.

From δ -wiggly permutations to δ -permutations. For all $j \in [n]$, erase $2j - 1$ and replace $2j$ by j if $\delta_j = +$, and erase $2j$ and replace $2j - 1$ by j if $\delta_j = -$.

From δ -permutations to δ -triangulations. Already described in [Rea06]. Given a permutation π , construct the union of the upper hulls of the point sets $[n]^- \triangle \pi([j])$ for all $j \in \{0, \dots, n\}$. \square

Recall that the δ -Cambrian lattice can be realized by the δ -Cambrian fan [RS09] and the δ -associahedron of [HL07]. Namely, to each diagonal (i, j) of the δ -gon, we associate the subset $X(\delta)$ of points of the δ -gon distinct from 0 and $n + 1$, and located below the line (i, j) , including i if and only if $\delta_i = +$ and j if and only if $\delta_j = +$. The δ -Cambrian fan is the fan with a ray $\mathbf{g}(d) := \pi(\mathbf{1}_{X(d)})$ for each internal diagonal d of the δ -gon and a cone $\mathbb{R}_{\geq 0} \langle \mathbf{g}(d) \mid d \in D^\circ \rangle$ for each dissection D of the δ -gon. The δ -associahedron Asso_δ is the polytope defined by the equality $\langle \mathbf{1}_n \mid \mathbf{x} \rangle = \binom{n+1}{2}$ and the inequalities $\langle \mathbf{g}(d) \mid \mathbf{x} \rangle \leq \binom{|X(d)|+1}{2}$ for all internal diagonals d of the δ -gon.

Proposition 73. *The δ -associahedron Asso_δ is normally equivalent to the face of the wigglyhedron \mathbb{W}_n corresponding to the wiggly pseudodissection formed by the δ -wiggly arcs.*

Proof. Following the proof of Proposition 72, consider the map ζ sending a diagonal (i, j) of the δ -gon to the wiggly arc $(0, j, [i] \cup]i, j[^- ,]i, j[^+)$ if $\delta_i \geq 0$ and $(0, j,]i, j[^-, [i] \cup]i, j[^+)$ if $\delta_i \leq 0$. Let $a := (i, j)$ and $a' := (i', j')$ be two crossing diagonals of the δ -gon. Then any δ -triangulations T and T' with $T \setminus \{a\} = T' \setminus \{a'\}$ contain the diagonals $b := (i', j)$ and $b' := (i, j')$ of the δ -gon, and $\mathbf{g}(a) + \mathbf{g}(a') = \mathbf{g}(b) + \mathbf{g}(b')$. Moreover, if $\alpha := \zeta(a)$, $\alpha' := \zeta(a')$, $\beta := \zeta(b)$ and $\beta' := \zeta(b')$, then $\mathbf{g}(\alpha) + \mathbf{g}(\alpha') = \mathbf{g}(\beta) + \mathbf{g}(\beta')$. We thus conclude that ζ preserves the linear dependences among \mathbf{g} -vectors. Hence, there is a linear map $Z : \mathbb{H}_n \rightarrow \mathbb{H}_{2n}$ such that $Z(\mathbf{g}(d)) = \mathbf{g}(\zeta(d))$ for all diagonals d of the δ -gon. This map is a normal equivalence from the δ -associahedron Asso_δ to the face of the wigglyhedron \mathbb{W}_n corresponding to the wiggly pseudodissection formed by the δ -wiggly arcs. \square

Remark 74. In their work on polytopal realizations of \mathbf{g} -vector fans of finite type cluster algebras [HPS18], C. Hohlweg, V. Pilaud and S. Stella already constructed a *universal associahedron* \mathbb{U}_n which somehow contains all Cambrian fans (actually all \mathbf{g} -vector fans from any initial cluster seed). More precisely, the δ -Cambrian fan for each $\delta \in \{+, -\}^n$ appears as a section of the normal fan of the universal associahedron \mathbb{U}_n , which is a $(n^2 + n - 4)/2$ -dimensional polytope. In contrast, Proposition 73 states that the δ -Cambrian fan for each $\delta \in \{+, -\}^n$ appears as a link of the normal fan of the wigglyhedron \mathbb{W}_n , which is a $(2n - 1)$ -dimensional polytope. There seem to be no connection between these two universal polytopes. However, this certainly raises the question to construct analogues of the wigglyhedron for all finite Coxeter groups.

4.2. **Wiggly pseudotriangulations of planar point sets.** We now consider Theorem 63 and Conjecture 44 for wiggly pseudotriangulations of arbitrary point sets in the plane (neither necessarily aligned, nor necessarily in general position).

Definition 75. Fix a point set P of the plane, and an arbitrary total order $<$ on P . A *wiggly arc* is a quadruple (p, q, R, S) where $p < q \in P$ and the sets R and S form a partition of the points of P located in the open segment joining p to q . Two wiggly arcs (p, q, R, S) and (p', q', R', S') are *crossing* if

- either the segments $[p, q]$ and $[p', q']$ cross,
- or $(R \cap S') \cup (\{p, q\} \cap S') \cup (R \cap \{p', q'\}) \neq \emptyset \neq (R' \cap S) \cup (\{p', q'\} \cap S) \cup (R' \cap \{p, q\})$,

A set X of wiggly arcs is *pointed* if for any $p \in P$, the wiggly arcs of X with an endpoint at p generate a pointed cone. The *wiggly complex* WC_P is the simplicial complex of pairwise pointed and non-crossing subsets of wiggly arcs. Note that the boundary wiggly arcs are irrelevant, which allows us to consider a reduced wiggly complex WC_P induced by internal wiggly arcs. A *wiggly pseudotriangulation* of P is a facet of WC_P . The *wiggly flip graph* WFG_P is the adjacency graph of the facets of WC_P . See Figure 15 for an illustration.




FIGURE 15. The wiggly complex WC_P (left) and the wiggly flip graph WFG_P (right) of a point set P .

Note that Definition 75 recovers two specific situations, namely

- Definitions 17 and 21 in the case of aligned points, and
- the pseudodissection complex of P and pseudotriangulation flip graph of P in the case of points P in general position. We refer to the original article of M. Pocchiola and G. Vegter [PV96] and to the nice survey of G. Rote, F. Santos and I. Streinu [RSS08].

Observe that it is not even clear from Definition 75 that the wiggly complex WC_P is a pure pseudomanifold (hence that the wiggly flip graph WFG_P is well-defined). However, we make the following ambitious conjecture.

Conjecture 76. *For any point set P in the plane, the wiggly complex WC_P is the boundary complex of a simplicial polytope.*

In Conjecture 76, the case of aligned points is given by the wigglyhedron of Theorem 63, while the case of points in general position is given by the pseudotriangulation polytope of [RSS03]. This raises in particular the following question.

Question 77. Can the construction of Theorem 63 be adapted to provide a more combinatorial construction of the polytope of pseudotriangulations [RSS03], that would only depend on the order type (*a.k.a.* oriented matroid [BLS⁺99]) of the point configuration?

Using the duality between lines in the plane and points in the Möbius strip, the pseudotriangulations of [PV96, RSS08] were interpreted in [PP12] as pseudoline arrangements on sorting networks. It is natural to look for the analogue of this interpretation in the wiggly setting.

Question 78. Is there a dual interpretation of wiggly pseudotriangulations as some sort of pseudoline arrangements?

This dual interpretation enables [PP12] to consider the pseudotriangulations of [PV96, RSS08] and the multistriangulations of [PS09] under the same roof, and to define multi-pseudo-triangulations. The extension to the wiggly case is a natural question.

Question 79. Is there a multi wiggly complex?

Moreover, pseudoline arrangements on sorting networks can also be interpreted as (type A) subword complexes of [KM04], which in turn extend to arbitrary finite Coxeter group. See [Stu11, CLS14] for the study of subword complexes corresponding to cluster complexes and multi-cluster complexes.

Question 80. Can the wiggly complex be extended to arbitrary finite Coxeter groups?

Finally, it is natural to consider the extension of Conjecture 44.

Question 81. Is the wiggly flip graph WFG_P Hamiltonian for any point set P in the plane?

5. CATEGORICAL INTERPRETATION OF WIGGLY PSEUDOTRIANGULATIONS

The aim of this section is to explain a categorical interpretation of wiggly arcs. We first present a particular triangulated category (Section 5.1), and recall results of Khovanov–Seidel [KS02] (Section 5.2) that interpret the objects of this category as wiggly arcs, such that intersections between wiggly arcs correspond to morphisms in this category. We then explain how wiggly pseudotriangulations govern the combinatorial structure of a natural categorical decomposition of objects (Section 5.3). We also discuss categorical conditions for two wiggly arcs to be compatible as in Definition 3 (Section 5.4).

5.1. A triangulated category associated to a quiver. We begin by describing the category of interest, which is a subcategory of $D^b(A_m\text{-mod})$ for a certain finite-dimensional algebra A_m . That is, a subcategory of the bounded derived category of finitely generated A_m -modules. The algebra A_m , called the *zigzag algebra*, is defined as a quotient of the path algebra of the quiver Q_m with m vertices depicted in Figure 16. Let kQ_m be the path algebra of Q_m , graded by path length. Let e_i denote the idempotent path at the vertex i , and let (i, j) denote the arrow from vertex i to a (neighbouring) vertex j . A longer path, which is a composition of arrows, will be denoted by (i_1, i_2, \dots, i_n) .

Definition 82. The *zigzag algebra* A_m corresponding to the quiver Q_m depicted in Figure 16 is the quotient of kQ_m by the relations $(i - 1, i, i + 1) = 0 = (i + 1, i, i - 1)$ and $(i, i - 1, i) = (i, i + 1, i)$ for all $1 < i < m$.



FIGURE 16. The quiver Q_m . The algebra A_m is a quotient of the path algebra kQ_m .

Remark 83. The definition of the zigzag algebra as used here originally appeared in [HK01, Sect. 3]. This is the “un-signed” zigzag algebra. It is technically better to use a “signed” zigzag algebra, and the comparison between the two is discussed in [BDL20, Rem. 6.6]. As the two versions are isomorphic in our example, we ignore this technicality and use the easier definition.

Let P_i denote the indecomposable projective A_m -module defined as $P_i = A_m \cdot e_i$. For any integer j , let $P_i\langle j \rangle$ denote its graded shift by j . There is an embedding of A_m -mod into $D^b(A_m\text{-mod})$, which sends any A_m -module as a complex of A_m -modules that is concentrated in degree 0. In particular, the objects $P_i\langle j \rangle$ can be regarded as objects of $D^b(A_m\text{-mod})$.

Definition 84. The category \mathcal{C}_m is defined to be the smallest triangulated subcategory of $D^b(A_m\text{-mod})$ containing all the objects $P_i\langle j \rangle$.

For further discussion on the construction, see *e.g.* [BDL22, Sect. 2.3.3]. The category \mathcal{C}_m is k -linear and triangulated, where $[1]$ denotes the triangulated shift. For $X, Y \in \mathcal{C}_m$, let $\text{Hom}^{g,h}(X, Y)$ denote the space of morphisms in \mathcal{C}_m of internal degree g and homological degree h . In other words, we have

$$\text{Hom}^{g,h}(X, Y) = \text{Hom}(X, Y\langle g \rangle[h]).$$

Furthermore, for any integers a and b , we have

$$\text{Hom}^{g,h}(X\langle g' \rangle[h'], Y\langle g' \rangle[h']) = \text{Hom}^{g,h}(X, Y).$$

Remark 85. The generating objects P_i (and hence their graded and homological shifts) satisfy $\text{Hom}^{0,0}(P_i, P_i) = \text{Hom}^{2,0}(P_i, P_i) = \text{Hom}^{1,0}(P_i, P_{i\pm 1}) = k$ and $\text{Hom}^{g,h}(P_i, P_j) = 0$ in all other cases.

We now describe the t -structure induced by the full subcategory of *linear complexes* of \mathcal{C}_m . A variant of this construction is also described in [LQ21, Sect. 2.5]. It is convenient to think of an object P^\bullet of \mathcal{C}_m as a bigraded (projective) A_m -module equipped with a differential $d: P^\bullet \rightarrow P^\bullet$. More precisely, we can write

$$P^\bullet = \bigoplus_{g,h,i} M_i\langle g \rangle[h],$$

where g is the internal grading shift, h is the homological grading shift, and each object M_i is a direct sum of copies of objects in $\{P_1, \dots, P_m\}$. Set P^h to be the direct sum of all summands that have homological grading shift h . That is, in the notation above, we have

$$P^h = \bigoplus_{g,i} M_i\langle g \rangle[h].$$

Warning 86. By convention, if M_i is an A_m -module, then the object $M_i[h]$ is placed in homological degree $-h$. Since the differential d raises the homological degree by one, it lowers the homological shift by one. That is, we have

$$d: M_i\langle g \rangle[h] \rightarrow \bigoplus_{g',i'} M_{i'}\langle g' \rangle[h-1].$$

In other words, d maps P^h to P^{h-1} .

Definition 87. Let P^\bullet be an object of \mathcal{C}_m , described as a bigraded module

$$P^\bullet = \bigoplus M_i\langle g \rangle[h]$$

together with an internal differential $d: P^\bullet \rightarrow P^\bullet$ as above. We say that P^\bullet has *level* ℓ for some $\ell \in \mathbb{Z}$ if for each bigraded piece $M_i\langle g \rangle[h]$ of P^\bullet , we have $g + h = \ell$. We say that P^\bullet is *linear* if P^\bullet has level $\ell = 0$.

For any integer N , let $\mathcal{C}_m^{\leq N} \subset \mathcal{C}_m$ be the smallest full subcategory of \mathcal{C}_m closed under isomorphism, containing all complexes of level $\geq -N$. For any integer N , let $\mathcal{C}_m^{\geq N} \subset \mathcal{C}_m$ be the smallest full subcategory of \mathcal{C}_m closed under isomorphism, containing all complexes of level $\leq -N$.

We refer the reader to *e.g.* [BBD82, Sect. 1.3] for the definition of a t -structure on a triangulated category. The following proposition is easily checked using the calculations in Remark 85 and standard techniques involving categories of complexes; we omit the proof.

Proposition 88. *The pair of subcategories $(\mathcal{C}_m^{\leq 0}, \mathcal{C}_m^{\geq 0})$ forms a bounded t -structure on \mathcal{C}_m .*

We call this t -structure the *standard t -structure* on \mathcal{C}_m . Set $\mathcal{A}_m := \mathcal{C}_m^{\leq 0} \cap \mathcal{C}_m^{\geq 0}$ to be the heart of this t -structure; this is the smallest full subcategory of \mathcal{C}_m closed under isomorphism, which contains all linear complexes. It follows that \mathcal{A}_m is abelian.

Also record the following statements for future use; the first may be found in [BBD82, Sect. 1.3]

Definition/Proposition 89. Let $(\mathcal{C}^{\leq 0}, \mathcal{C}^{\geq 0})$ be a t -structure on a triangulated category \mathcal{C} with heart $\mathcal{A} := \mathcal{C}^{\leq 0} \cap \mathcal{C}^{\geq 0}$. For each integer i , we have the i th truncation functors

$$\tau^{\leq i}: \mathcal{C} \rightarrow \mathcal{C}^{\leq i} \quad \text{and} \quad \tau^{\geq i}: \mathcal{C} \rightarrow \mathcal{C}^{\geq i},$$

which are right and left adjoint respectively to the inclusion functors $\mathcal{C}^{\leq 0} \rightarrow \mathcal{C}$ and $\mathcal{C}^{\geq 0} \rightarrow \mathcal{C}$. The i th *cohomology functor* is defined to be the composition $\tau^{\leq i} \circ \tau^{\geq i}$. It is canonically isomorphic to the composition $\tau^{\geq i} \circ \tau^{\leq i}$, and maps an object of \mathcal{C} to an object of \mathcal{A} .

The next statement, easy to check from the definitions, is mentioned *e.g.* in [Bay11, Sect. 3.3].

Definition/Proposition 90. Let \mathcal{C} be any triangulated category, and $\mathcal{A} \subset \mathcal{C}$ the heart of a bounded t -structure on \mathcal{C} . Let X be a non-zero object of \mathcal{C} , and let k and ℓ be the maximum and minimum integers respectively for which $H^k(X)$ and $H^\ell(X)$ are non-zero. Then X has a unique *cohomology filtration*, which is the unique filtration of the form

$$\begin{array}{ccccccc} 0 & \longrightarrow & X_{-\ell} & \longrightarrow & \cdots & \longrightarrow & X_{-k+1} & \longrightarrow & X_{-k} = X \\ & & \swarrow & & \swarrow & & \swarrow & & \swarrow \\ & & H^\ell(X)[- \ell] & & H^{\ell+1}(X)[- \ell - 1] & & \cdots & & H^k(X)[- k] \end{array}$$

5.2. The Khovanov–Seidel curve model. In this section we briefly recall how to recover an object of \mathcal{C}_{n+1} from a wiggly arc on $n+2$ points on a line. In fact, the procedure works on more general curves as well (namely, those that are not necessarily x -monotone).

Consider $n+2$ points arranged in a straight horizontal line, and labelled $0, 1, \dots, n+1$ as in Section 2. In addition to the wiggly arcs, consider isotopy classes of curves whose endpoints lie on distinct marked points, and which do not cross themselves or any marked point in their interior. For convenience, we refer to such an isotopy class as a *curve*.

The Khovanov–Seidel correspondence (explained in Section 4 of [KS02]) associates objects of \mathcal{C}_{n+1} (up to an overall grading and homological shift) to curves on a configuration of $n+2$ marked points arranged in a straight line. We illustrate this correspondence and Proposition 91 in Figure 17. In particular, the line segment joining point i to point $i+1$ is sent to the object P_{i+1} in the category \mathcal{C}_{n+1} . Moreover, the minimal intersection number between any pair of curves is the number of morphisms between the associated objects [KS02, Prop. 4.9]. In fact, the proposition in [KS02] is a stronger result: by considering *bigraded curves*, we obtain *bigraded intersection numbers* between the two associated objects.

Any object X obtained via the Khovanov–Seidel correspondence is *spherical*; that is, $\text{Hom}^{g,0}(X, X)$ is k when $g = 0, 2$, and $\text{Hom}^{g,h}(X, X) = 0$ in all other cases. A spherical object X is necessarily indecomposable as $\text{Hom}^{0,0}(X, X) = k$.

Proposition 91. *Let α be any curve, and let $X = X^\bullet$ be a chain complex that corresponds to α under the Khovanov–Seidel reconstruction. Recall that for any integer j , the cohomology object $H^j(X)$ is an object of \mathcal{A}_m , and hence a direct sum of indecomposable objects of \mathcal{A}_m .*

- (1) *The object X is a (spherical) object of \mathcal{A}_m up to shift if and only if α is a wiggly arc. In this case, $H^j(X) \neq 0$ for exactly one j , such that $X \cong H^j(X)[-j]$.*




FIGURE 17. An illustration of the Khovanov–Seidel curve model and Proposition 91.

(2) If α is a general curve, each indecomposable summand of $H^j(X)$ for any j is a spherical object of \mathcal{A}_m that arises from a wiggly arc, up to shift.

Proof. Let X be a complex built from an arbitrary (not necessarily wiggly) curve α . Recall from [KS02] that X is spherical, and hence indecomposable. Let us now analyse the indecomposable summands of its cohomology objects.

We recall that the procedure to build up X from α explicitly describes the differential. We analyse the possible non-zero components of the differential, checking how each interacts with the level. The only two possibilities are as follows:

- (1) a map $P_i\langle g\rangle[h] \rightarrow P_{i\pm 1}\langle g+1\rangle[h-1]$, namely multiplication on the right by the path $(i, i \pm 1)$, which does not change the level; or
- (2) a map $P_i\langle g\rangle[h] \rightarrow P_i\langle g+2\rangle[h-1]$, namely multiplication on the right by the “loop” path $(i, i \pm 1, i)$, which raises the level by one.

The first possibility appears each time the curve wiggles either above the i th point or below the $(i-1)$ th point. The second possibility appears each time the curve makes a U-turn around the right of i th or the left of the $(i-1)$ th point.

Computing the j th cohomology $H^j(X)$ simply amounts to truncating the complex to level j . Explicitly, the truncation is a new chain complex in which we only keep the terms of X that have level j , together with all maps between them, and forget all the other terms. From the calculation above, we see that the resulting complex only has components in its differential of the first type, namely $P_i\langle g\rangle[h] \rightarrow P_{i\pm 1}\langle g+1\rangle[h-1]$ where $g+h=j$. We kill all components of the second type (the “loop” components). The resulting chain complex breaks up into indecomposable summands, each of which is an object of $\mathcal{A}_m[j]$, and this direct sum is $H^j(X)[j]$.

The object X is in the heart up to shift if and only if it is concentrated in a single level j . Truncating to level j as described above, we do not delete any of the terms of X . This can happen if and only if the differential of X contains no “loop” components, which appear in the complex if and only if α makes a U-turn around one of the $n+2$ points. Thus X is in the heart up to shift if and only if α is x -monotone. This proves the first part of the proposition.

More generally, consider the indecomposable pieces of the level- j truncations of X for various j , up to shift. As explained above, in terms of the curve α , this corresponds to breaking up the curve α into x -monotone portions. More precisely, we begin at an endpoint p_0 of α and follow the curve until it either ends at a point p_1 , or stops being x -monotone by making a U-turn around a

point p_1 . In this case, we take the portion of α between p_0 and p_1 , calling it α_1 , and setting its endpoints to be p_0 and p_1 respectively. Note that $p_0 \neq p_1$ because α is x -monotone between p_0 and p_1 . We continue with the remainder of α starting at p_1 , greedily taking the maximal x -monotone portion each time, to obtain pieces $\alpha_1, \alpha_2, \dots, \alpha_k$, which are all wiggly arcs. These wiggly arcs correspond to complexes X_1, \dots, X_k , which are spherical objects of \mathcal{A}_m , up to shift. It is clear from the construction of the associated complex that these objects, up to shift, are exactly the indecomposable pieces of the cohomology truncations $H^j(X)$ for various j . This argument proves the second part of the proposition. \square

5.3. Cohomology filtrations of spherical objects. In this section we explain how wiggly pseudotriangulations govern the combinatorial structure of a natural categorical decomposition of spherical objects of \mathcal{C}_m .

For a general object X of \mathcal{C}_m , the indecomposable factors of the cohomology filtration can be arbitrary. However, it turns out that the cohomology filtration is highly constrained if X is spherical, and these constraints are given by wiggly pseudotriangulations.

Proposition 92. *Let $X \in \mathcal{C}_{n+1}$ be a spherical object. Consider its cohomology filtration*

$$\begin{array}{ccccccc}
 0 & \xrightarrow{\quad} & X_{-\ell} & \xrightarrow{\quad} & \cdots & \xrightarrow{\quad} & X_{-k} = X \\
 & \swarrow & \searrow & \swarrow & \searrow & \swarrow & \searrow \\
 & & H^\ell(X)[- \ell] & & H^{\ell+1}(X)[- \ell - 1] & & \cdots & & H^k(X)[- k]
 \end{array}$$

Consider the set of all possible indecomposable summands of the cohomology objects $\{H^j(X)\}$ taken without multiplicity. Let T be corresponding set of wiggly arcs, as explained in Proposition 91. Then

- (1) T is a subset of a wiggly pseudotriangulation,
- (2) T contains at most one of the two irrelevant wiggly arcs.

Proof. The proof of this proposition is an easy consequence of analysing the cohomology filtration from the point of view of curves on the disk with marked points. Let X_1, \dots, X_k be the objects in T without repetition, which arise from wiggly arcs $\alpha_1, \dots, \alpha_k$ as explained in the proof of Proposition 91. Let $\beta = (i, j, A, B)$ and $\beta' = (i', j', A', B')$ be two distinct wiggly arcs among $\{\alpha_1, \dots, \alpha_k\}$. We check that β and β' are compatible in the sense of Definition 3.

Recall that β and β' are two pieces of the curve α . Therefore β and β' cannot cross, because α does not cross itself. Suppose that β and β' are non pointed, and without loss of generality, suppose that $i' = j$. This means that the piece β was cut from α either because $i' = j$ is an endpoint of α , or because α loops around the right of the point i' . Moreover, if i' is an endpoint of α , then the portion of α closest to i' lies to the left of i' .

Similarly, the piece β' was cut from α either because $i' = j$ is an endpoint of α , or because α loops around the left of the point j . Moreover, if j is an endpoint of α , then the portion of α closest to j lies to the right of j .

We have four possible combinations:

- (1) $i' = j$ is an endpoint such that α travels left starting at i' , and $i' = j$ is an endpoint such that α travels right starting at j ;
- (2) $i' = j$ is an endpoint such that α travels left starting at i' , and α loops around the left of the point j ;
- (3) α loops around the right of the point $i' = j$, and $i' = j$ is an endpoint such that α travels right starting at j ;
- (4) α loops around the right of the point $i' = j$, and α loops around the left of the point j .

It is clear that none of these possibilities can occur within the curve α , and hence β and β' must be pointed. Therefore $T = \{\alpha_1, \dots, \alpha_k\}$ is a subset of a wiggly pseudotriangulation.

Now suppose for contradiction that T contains both external edges. Then some portion of α curves around each of the extreme endpoints 0 and $n + 1$. Consider the outermost strands of α that curve around 0 and $n + 1$ respectively. These must then connect to each other, resulting in α having a separate closed component that winds around the outside of the $n + 2$ points. This is

not possible because α is a connected curve with two endpoints. Therefore T cannot contain both external edges. \square

Remark 93. Note that in Definition 21, we adopted the convention that the faces of the wiggly complex contain none of the two external wiggly arcs. In contrast, the wiggly dissections of Proposition 92 contain at most one of the two external wiggly arcs. Combinatorially, this is just the Cartesian product of the wiggly complex with an edge.

5.4. Categorical constraints on compatibility of curves. In this section, we give a categorical criterion for two wiggly arcs to be compatible as in Definition 3. As such, this section is somewhat dual to the previous section: the criterion in this section determines whether a pair of wiggly arcs can belong together in a single wiggly pseudotriangulation, whereas the previous section generated a wiggly pseudotriangulation given any object of the category.

Recall that two wiggly arcs α and α' are said to be compatible (as in Definition 3) if they are non-crossing and pointed. The incompatibility degree of α and α' (as in Definition 59) is 1 if α and α' are non pointed, and the number of crossings of α and α' otherwise.

Proposition 94. *Let X and X' be objects in the standard heart \mathcal{A}_m corresponding to two wiggly arcs α and α' respectively. Then α and α' are compatible if and only if $\text{Ext}^1(X, X') = 0$. More generally, the incompatibility degree of α and α' is the dimension of $\text{Ext}^1(X, X')$.*

Proof. The proof follows from a specialisation of the bigraded intersection number calculations that appear in Sections 3 and 4 of [KS02]. The bigraded intersection number as defined in [KS02, Sect. 3d] between the curves α and α' is a polynomial in the variables q_1^\pm, q_2^\pm . It follows from [KS02, Prop. 4.9] that this bigraded intersection number equals the Poincaré polynomial of the morphisms between X and X' , namely the expression

$$(1) \quad \sum_{s_1, s_2} \dim(\text{Hom}_{\mathcal{C}_m}(X, X' \langle -s_2 \rangle [s_1])) q_1^{s_1} q_2^{s_2}.$$

However, the grading convention in [KS02] is not the same as ours — our grading is by path length, while the grading in [KS02] is one for arrows in one direction and zero for arrows in the opposite direction. So we have to modify the gradings when we construct the complexes. In particular, the table in [KS02, Lem. 3.20] that computes the bigraded intersection numbers will change, and so will the table in [KS02, Lem. 4.12] that computes the Poincaré polynomials of $\text{Hom}(P_k, -)$.

With our path-length grading, note that a morphism in $\text{Hom}_{\mathcal{C}_m}(X, X' \langle -s_2 \rangle [s_1])$ has level $s_1 - s_2$; that is, it takes an object X of level 0 to the object $X' \langle -s_2 \rangle [s_1]$ of level $s_1 - s_2$. We can thus obtain the dimension of all morphisms of level ℓ by setting $q_1 = q$ and $q_2 = q^{-1}$ and taking the coefficient of q^ℓ in the expression (1). The dimension of $\text{Ext}^1(X, X')$, is therefore the coefficient of q under this specialisation.

It can be checked that under the path-length grading, the space of morphisms arising from an intersection of curves of “crossing” type as well as “non pointed” type contains exactly one morphism (up to equivalence) of level one; that is, a unique element of Ext^1 . Morphisms arising from other intersections do not contribute to Ext^1 . This calculation proves the proposition. \square

Remark 95. Proposition 94 is very similar to the condition of being compatible in a cluster-tilting object of a cluster category (see, e.g. [BMR⁺06, Cor. 4.3]) or in a τ -tilting object (see, e.g. [AIR14]). The condition in terms of crossings of curves already appears for some specializations (see, e.g. [FST08, PPP19]). However, in Proposition 94, it depends on shifting the objects X and X' so that they both lie in the standard heart, which is an auxiliary step that is not required in cluster-tilting and τ -tilting theory.

Remark 96. As stated, Proposition 94 in combination with Proposition 92 is not expected to be true in greater generality. For instance, for the zigzag algebra of Dynkin type D_4 , one can find two spherical objects with a non-trivial Ext^1 that nevertheless appear together in the cohomology filtration of another spherical object. However, we expect to see similar constraints on compatibility, which we hope to turn to in forthcoming work.

Recall from Remark 60 that two wiggly arcs α and α' are exchangeable if and only if their incompatibility degree is one. Recall from Proposition 20 the wiggly arcs β and β' associated to α and α' . Finally, recall from Lemma 55 that the \mathbf{g} -vectors of the wiggly arcs α , α' , β , and β' satisfy a linear dependence, depending on their type. Let us interpret this dependence categorically.

Let X , X' , Y and Y' be spherical objects in the standard heart that correspond to α , α' , β and β' respectively. First, note by Proposition 94 that X and X' are exchangeable if and only if there is a unique non-trivial extension between the corresponding objects of the standard heart. Next, let us understand Y and Y' categorically.

Proposition 97. *For α , α' , β , β' , and X , X' , Y , Y' as just discussed,*

- if α and α' are not pointed, then we have extensions

$$0 \rightarrow X' \rightarrow Y \rightarrow X \rightarrow 0 \quad \text{and} \quad 0 \rightarrow X \rightarrow Y' \rightarrow X' \rightarrow 0,$$

- if α and α' are crossing, then we have an extension

$$0 \rightarrow X' \rightarrow (Y \oplus Y') \rightarrow X \rightarrow 0.$$

Proof. The proof is once again a consequence of analysing the extensions explicitly via the complexes from the Khovanov–Seidel reconstruction. We do not reproduce the proof here, as the argument is contained in the proof of [Tho06, Thm. 5.5]. (See also [Tho06, Fig. 4].) \square

Note that if we write down the relations in the Grothendieck group of our category that correspond to the extensions from each case of Proposition 97, then we obtain precisely the linear dependences that appear between the \mathbf{g} -vectors in Lemma 55. More precisely, we have that

- if α and α' are not pointed, then $[Y] = [X] + [X'] = [Y']$, from which we see $[X] + [X'] = ([Y] + [Y'])/2$;
- if α and α' are crossing, then $[X] + [X'] = [Y] + [Y']$.

This suggests that there should be a way to construct the \mathbf{g} -vectors categorically in such a way that the \mathbf{g} -vectors also map to the Grothendieck group of the category. The \mathbf{g} -vector of an object X corresponding to a wiggly arc α can, of course, be reconstructed from α . However, at the moment we do not have a purely categorical construction of the \mathbf{g} -vector of X that does not refer to α , and it would be interesting to find this.

Question 98. Suppose we have a collection of spherical objects in the standard heart \mathcal{A}_m that satisfy the compatibility condition of Proposition 94. That is, the Ext^1 between any pair of objects in this collection is zero.

- (1) Can we prove properties about a maximal such collection without reference to curves?
- (2) In particular, can we understand the combinatorics of maximal such collections without reference to curves?
- (3) Given a maximal compatible collection of objects, can we always find a larger spherical object such that the factors of the cohomology filtration of that object are precisely shifts of the objects in the given collection?

ACKNOWLEDGEMENTS

This work started during the *Winterbraids XIII: School on low dimensional topology, braid groups and interactions*. We are grateful to the organizers B er enice Delcroix-Oger, Peter Feller, Vincent Florens, Hoel Queffelec for this fruitful event. We thank Anand Deopurkar and Anthony M. Licata for many useful conversations and ideas.

REFERENCES

- [AIR14] Takahide Adachi, Osamu Iyama, and Idun Reiten. τ -tilting theory. *Compos. Math.*, 150(3):415–452, 2014.
- [Bay11] Arend Bayer. A tour to stability conditions on derived categories. 2011.
- [BBD82] Alexander A. Beilinson, Joseph Bernstein, and Pierre Deligne. Faisceaux pervers. In *Analysis and topology on singular spaces, I (Luminy, 1981)*, volume 100 of *Ast erisque*, pages 5–171. Soc. Math. France, Paris, 1982.

- [BDL20] Asilata Bapat, Anand Deopurkar, and Anthony M. Licata. A Thurston compactification of the space of stability conditions. 2020. <http://arxiv.org/abs/2011.07908>.
- [BDL22] Asilata Bapat, Anand Deopurkar, and Anthony M. Licata. Spherical objects and stability conditions on 2-Calabi–Yau quiver categories. *Mathematische Zeitschrift*, 303(1), Dec 2022.
- [BLS⁺99] Anders Björner, Michel Las Vergnas, Bernd Sturmfels, Neil White, and Günter M. Ziegler. *Oriented matroids*, volume 46 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, Cambridge, second edition, 1999.
- [BMR⁺06] Aslak Bakke Buan, Bethany Marsh, Markus Reineke, Idun Reiten, and Gordana Todorov. Tilting theory and cluster combinatorics. *Advances in Mathematics*, 204(2):572–618, Aug 2006.
- [BR87] M. C. R. Butler and Claus Michael Ringel. Auslander-Reiten sequences with few middle terms and applications to string algebras. *Comm. Algebra*, 15(1-2):145–179, 1987.
- [CFZ02] Frédéric Chapoton, Sergey Fomin, and Andrei Zelevinsky. Polytopal realizations of generalized associahedra. *Canad. Math. Bull.*, 45(4):537–566, 2002.
- [CLS14] Cesar Ceballos, Jean-Philippe Labbé, and Christian Stump. Subword complexes, cluster complexes, and generalized multi-associahedra. *J. Algebraic Combin.*, 39(1):17–51, 2014.
- [CP16] Cesar Ceballos and Vincent Pilaud. The diameter of type D associahedra and the non-leaving-face property. *European J. Combin.*, 51:109–124, 2016.
- [DRS10] Jesus A. De Loera, Jörg Rambau, and Francisco Santos. *Triangulations: Structures for Algorithms and Applications*, volume 25 of *Algorithms and Computation in Mathematics*. Springer Verlag, 2010.
- [FST08] Sergey Fomin, Michael Shapiro, and Dylan Thurston. Cluster algebras and triangulated surfaces I. Cluster complexes. *Acta Math.*, 201(1):83–146, 2008.
- [FZ02] Sergey Fomin and Andrei Zelevinsky. Cluster algebras. I. Foundations. *J. Amer. Math. Soc.*, 15(2):497–529, 2002.
- [FZ03] Sergey Fomin and Andrei Zelevinsky. Cluster algebras. II. Finite type classification. *Invent. Math.*, 154(1):63–121, 2003.
- [HHMW22] Elizabeth Hartung, Hung P. Hoang, Torsten Mütze, and Aaron Williams. Combinatorial generation via permutation languages. I. Fundamentals. *Trans. Amer. Math. Soc.*, 375(4):2255–2291, 2022.
- [HK01] Ruth Stella Huerfano and Mikhail Khovanov. A category for the adjoint representation. *J. Algebra*, 246(2):514–542, 2001.
- [HL07] Christophe Hohlweg and Carsten Lange. Realizations of the associahedron and cyclohedron. *Discrete Comput. Geom.*, 37(4):517–543, 2007.
- [HLT11] Christophe Hohlweg, Carsten Lange, and Hugh Thomas. Permutahedra and generalized associahedra. *Adv. Math.*, 226(1):608–640, 2011.
- [HPS18] Christophe Hohlweg, Vincent Pilaud, and Salvatore Stella. Polytopal realizations of finite type \mathfrak{g} -vector fans. *Adv. Math.*, 328:713–749, 2018.
- [KM04] Allen Knutson and Ezra Miller. Subword complexes in Coxeter groups. *Adv. Math.*, 184(1):161–176, 2004.
- [KS02] Mikhail Khovanov and Paul Seidel. Quivers, Floer cohomology, and braid group actions. *J. Amer. Math. Soc.*, 15(1):203–271, 2002.
- [LQ21] Anthony M. Licata and Hoel Queffelec. Braid groups of type ade , garside monoids, and the categorified root lattice. *Annales Scientifiques de l’École Normale Supérieure*, 54(2):503–548, 2021.
- [Pou14] Lionel Pournin. The diameter of associahedra. *Adv. Math.*, 259:13–42, 2014.
- [PP12] Vincent Pilaud and Michel Pocchiola. Multitriangulations, pseudotriangulations and primitive sorting networks. *Discrete Comput. Geom.*, 48(1):142–191, 2012.
- [PPP19] Yann Palu, Vincent Pilaud, and Pierre-Guy Plamondon. Non-kissing and non-crossing complexes for locally gentle algebras. *J. Comb. Algebra*, 3(4):401–438, 2019.
- [PPP21] Yann Palu, Vincent Pilaud, and Pierre-Guy Plamondon. Non-kissing complexes and tau-tilting for gentle algebras. *Mem. Amer. Math. Soc.*, 274(1343), 2021.
- [PS09] Vincent Pilaud and Francisco Santos. Multitriangulations as complexes of star polygons. *Discrete Comput. Geom.*, 41(2):284–317, 2009.
- [PV96] Michel Pocchiola and Gert Vegter. Topologically sweeping visibility complexes via pseudotriangulations. *Discrete Comput. Geom.*, 16(4):419–453, 1996.
- [Rea06] Nathan Reading. Cambrian lattices. *Adv. Math.*, 205(2):313–353, 2006.
- [Rea15] Nathan Reading. Noncrossing arc diagrams and canonical join representations. *SIAM J. Discrete Math.*, 29(2):736–750, 2015.
- [RS09] Nathan Reading and David E. Speyer. Cambrian fans. *J. Eur. Math. Soc.*, 11(2):407–447, 2009.
- [RSS03] Günter Rote, Francisco Santos, and Ileana Streinu. Expansive motions and the polytope of pointed pseudo-triangulations. In *Discrete and computational geometry*, volume 25 of *Algorithms Combin.*, pages 699–736. Springer, Berlin, 2003.
- [RSS08] Günter Rote, Francisco Santos, and Ileana Streinu. Pseudo-triangulations—a survey. In *Surveys on discrete and computational geometry*, volume 453 of *Contemp. Math.*, pages 343–410. Amer. Math. Soc., Providence, RI, 2008.
- [San12] Francisco Santos. A counterexample to the Hirsch conjecture. *Ann. of Math. (2)*, 176(1):383–412, 2012.

- [San13] Francisco Santos. Recent progress on the combinatorial diameter of polytopes and simplicial complexes. *TOP*, 21(3):426–460, 2013.
- [Stu11] Christian Stump. A new perspective on k -triangulations. *J. Combin. Theory Ser. A*, 118(6):1794–1800, 2011.
- [Tho06] Richard P. Thomas. Stability conditions and the braid group. *Commun. Anal. Geom.*, 14(1):135–161, 2006.
- [Zie98] Günter M. Ziegler. *Lectures on Polytopes*, volume 152 of *Graduate texts in Mathematics*. Springer-Verlag, New York, 1998.

THE AUSTRALIAN NATIONAL UNIVERSITY
Email address: `asilata.bapat@anu.edu.au`
URL: <https://asilata.github.io>

UNIVERSITAT DE BARCELONA
Email address: `vincent.pilaud@ub.edu`
URL: <https://www.ub.edu/comb/vincentpilaud/>